Pain after laparoscopic bilateral hernioplasty

Early results of a prospective randomized double-blind study comparing fibrin versus staples

E. Boldo · Andres Armelles · Guillermo Perez de Lucia · Felix Martin ·
Juan Pablo Aracil · Juan Manuel Miralles · David Martinez ·
Javier Escrig

Published online: 13 February 2008
© Springer Science+Business Media, LLC 2008

Erratum to: Surg Endosc DOI 10.1007/s00464-007-9587-z

The following coauthors were inadvertently omitted:

Andres Armelles; Consorcio Hospitalario Provincial
Castellon, Surgery

Guillermo Perez de Lucia; Consorcio Hospitalario Pro-
vincial Castellon, Surgery

Felix Martin; Consorcio Hospitalario Provincial Cas-
tellon, Surgery

Juan Pablo Aracil; Consorcio Hospitalario Provincial
Castellon, Surgery

Juan Manuel Miralles

David Martinez

Javier Escrig

Springer regrets the error.

The online version of the original article can be found under doi:
[10.1007/s00464-007-9587-z](https://doi.org/10.1007/s00464-007-9587-z)

E. Boldo (✉) · A. Armelles · G. Perez de Lucia · F. Martin ·
J. P. Aracil · J. M. Miralles · D. Martinez · J. Escrig
Surgery, Consorcio Hospitalario Provincial Castellon, Castellon,
Castellon, Spain
e-mail: eboldo@yahoo.com