Optimization of sympathectomy to treat palmar hyperhidrosis

G. Bischof · A. E. P. Cameron · C. Connery ·
J. R. M. De Campos · M. Hashmonai ·
P. B. Licht · C. H. Schick

Received: 2 January 2013 / Accepted: 17 April 2013 / Published online: 10 May 2013
© Springer Science+Business Media New York 2013

To the Editor,

We read with interest the article by Deng et al. [1] and compliment them for the important review they performed. This review, which includes articles from the last decade, comes to a conclusion regarding the best level of sympathetic ablation for the treatment of primary palmar hyperhidrosis shared by many authors (T3). However, in a thorough review (246 references) on the subject comparing the correlation of the method for sympathetic ablation with the subsequent occurrence of compensatory sweating [2], the authors could not define any such correlation.

In reviewing your study, we found that five articles for the period 2000–2006 were not included. These articles report the ablation of single ganglia [3–7]. Were there other similar omissions for the years 2007–2010? We would appreciate the comment of the authors about these omissions. Would the inclusion of these series in their study possibly change their conclusion?

Disclosures G. Bischof, A. E. P. Cameron, C. Connery, J. R. M. De Campos, M. Hashmonai, P. B. Licht, and C. H. Schick have no conflicts of interest or financial ties to disclose.

References

1. Deng B, Tan Q-Y, Jiang Y-G et al (2011) Optimization of sympathectomy to treat palmar hyperhidrosis: the systematic review and meta-analysis of studies published during the past decade. *Surg Endosc* 25:1893–1901
2. Kopelman D, Hashmonai M (2008) The correlation between the method of sympathetic ablation for palmar hyperhidrosis and the occurrence of compensatory hyperhidrosis: a review. *World J Surg* 32:2353–2356
3. Singh B, Shaik AS, Moodley J et al (2002) Limited thoracoscopic ganglionectomy for primary hyperhidrosis. *S Afr J Surg* 40:50–53
4. Reisfeld R, Nguyen R, Pnini A (2000) Endoscopic thoracic sympathectomy for treatment of essential hyperhidrosis syndrome: experience with 650 patients. *Surg Laparosc Endosc Percutan Tech* 10:5–10
5. Doolabh N, Horswell S, Williams M et al (2004) Thoracoscopic sympathectomy for hyperhidrosis: indications and results. *Ann Thorac Surg* 77:410–414
6. Atkinson JLD, Fealey RD (2003) Sympathotomy instead of sympathectomy for palmar hyperhidrosis: minimizing postoperative compensatory hyperhidrosis. *Mayo Clin Proc* 78:167–172
7. Goh PMY, Cheah W-K, da Costa M et al (2000) Needleoscopic thoracic sympathectomy: treatment for palmar hyperhidrosis. *Ann Thorac Surg* 70:240–242

G. Bischof (✉)
Department of Surgery, St Josef Hospital, Vienna, Austria
e-mail: georg.bischof@sjk-wien.at

A. E. P. Cameron
Department of Surgery, The Ipswich Hospital, Ipswich, UK

C. Connery
Department of Surgery, St. Luke's-Roosevelt Hospital Center,
New York, NY, USA

J. R. M. De Campos
Department of Thoracic Surgery, University of Sao Paulo, São
Paulo, Brazil

M. Hashmonai
Faculty of Medicine, Technion—Israel Institute of Technology,
Kesalsaba, Israel

P. B. Licht
Department of Cardiothoracic Surgery, Odense University
Hospital, Odense, Denmark

C. H. Schick
German Hyperhidrosiscenter, Surgery Isar Clinics, Munich,
Germany