ERRATUM

Erratum to: Laparoscopic versus robot-assisted bilateral nerve-sparing radical prostatectomy: comparison of pentafecta rates for a single surgeon

Anastasios D. Asimakopoulos · Roberto Miano ·
Nicola Di Lorenzo · Enrico Spera ·
Giuseppe Vespasiani · Camille Mugnier

Published online: 31 August 2013
© Springer Science+Business Media New York 2013

Erratum to: Surg Endosc
DOI 10.1007/s00464-013-3046-9

In the Results section of the Abstract, the phrase:

“Of the 140 patients who failed pentafecta, 90 (64.3 %) missed a single parameter, and the difference between the groups was significant (80 % LRP vs 53.3 % RALP, $p = 0.007$).”

Should be changed as follows:

“Of the 140 patients who failed pentafecta, 90 (64.3 %) missed a single parameter. In these cases, erectile deficit was the leading cause of pentafecta failure, with a significant difference between groups (80 % LRP cases that missed potency recovery vs 53.3 % RALP, $p = 0.007$).”

The online version of the original article can be found under doi:[10.1007/s00464-013-3046-9](https://doi.org/10.1007/s00464-013-3046-9).

A. D. Asimakopoulos (✉) · E. Spera
UOC of Urology, Department of Surgery, University of Tor Vergata, Policlinico Casilino, Rome, Italy
e-mail: tasospao2003@yahoo.com

A. D. Asimakopoulos · C. Mugnier
Division of Urology, Clinique Saint-Augustin, Bordeaux, France

R. Miano · G. Vespasiani
UOC of Urology, Department of Surgery, University of Tor Vergata, Policlinico Tor Vergata, Rome, Italy

N. Di Lorenzo
Department of Surgical Sciences, University of Tor Vergata, Policlinico Tor Vergata, Rome, Italy