ERRATUM

Erratum to: Multivariate comparison of B-ultrasound guided and laparoscopic continuous circulatory hyperthermic intraperitoneal perfusion chemotherapy for malignant ascites

Ming-Chen Ba · Hui Long · Shu-Zhong Cui ·
Yun-Qiang Tang · Yin-Bing Wu · Xiang-Liang Zhang ·
Hong-Sheng Tang · Sai-Xi Bai

Published online: 24 January 2014
© Springer Science+Business Media New York 2014

Erratum to: Surg Endosc (2013) 27:2735–2743
DOI 10.1007/s00464-013-2800-3

The right data of the incidence of port-site metastasis in B ultrasound-guided group should be 9.375 %, in laparoscopically-assisted group should be 10 %, but the error data of our calculation in B ultrasound-guided group was 18.75 %, and in laparoscopically-assisted group was 18.18 % in text and table of the published paper.

The online version of the original article can be found under
doi:[10.1007/s00464-013-2800-3](https://doi.org/10.1007/s00464-013-2800-3).

M.-C. Ba (✉) · S.-Z. Cui · Y.-Q. Tang · Y.-B. Wu ·
X.-L. Zhang · H.-S. Tang · S.-X. Bai
Intracelom Hyperthermic Perfusion Therapy Center,
Cancer Hospital of Guangzhou Medical College,
Guangzhou 510095, China
e-mail: bamingchen2011@126.com

H. Long (✉)
Department of Pharmacy, Guangzhou Dermatology Institute,
Guangzhou 510095, China
e-mail: 798587414@qq.com