ERRATUM

Erratum to: Laparoscopic resection of paraaortic/paracaval neurogenic tumors: surgical outcomes and technical tips

Takashige Abe¹ · Ataru Sazawa² · Toru Harabayashi³ · Yuichiro Oishi¹ · Naoto Miyajima¹ · Kunihiro Tsuchiya¹ · Satoru Maruyama¹ · Hiromi Okada⁴ · Nobuo Shinohara¹

Published online: 7 July 2016
© Springer Science+Business Media New York 2016

Erratum to: Surg Endosc DOI 10.1007/s00464-015-4740-6

There were errors in Table 2 as originally published. Regarding the margin status, negative: $n = 19$ and positive: $n = 6$ were correct. The correct version of Table 2 is reproduced here.

Table 2 Patients' outcomes

	$n = 25$
Operative time (min)	Median 208 (range 73–513)
Blood loss (mL)	Median 10 (range 0–1020)
Intraoperative complication	
Injury of vena cava	1
Injury of duodenum	1
Postoperative complication	
Depression	1 (Grade 2)
Hypoglycemia	1 (Grade 2)
Chylous leakage	1 (Grade 1)
Postoperative hospital stay (days)	Median 8 (range 5–13)
Pathology	
Paraganglioma	12
Ganglioneuroma	7
Schwannoma	6
Margin	
Negative	19
Positive	6

The online version of the original article can be found under doi:[10.1007/s00464-015-4740-6](https://doi.org/10.1007/s00464-015-4740-6).

✉ Takashige Abe
takataka@rf6.so-net.ne.jp

¹ Department of Urology, Hokkaido University Graduate School of Medicine, North-15, West-7, North Ward, Sapporo 060-8638, Japan

² Department of Urology, Obihiro-Kosei General Hospital, Obihiro, Japan

³ Department of Urology, Hokkaido Cancer Center, Sapporo, Japan

⁴ Department of Surgical Pathology, Hokkaido University Hospital, Sapporo, Japan