Correction to: Comparison between submucosal tunneling endoscopic resection and video-assisted thoracoscopic enucleation for esophageal submucosal tumors originating from the muscularis propria layer: a randomized controlled trial

Ningli Chai¹ · Chen Du¹ · Ying Gao¹ · Xiaotong Niu¹ · Yaqi Zhai¹ · Enqiang Linghu¹ · Yang Liu² · Bo Yang² · Zhongsheng Lu¹ · Zhenjuan Li¹ · Xiangdong Wang¹ · Ping Tang¹

Published online: 23 March 2018
© Springer Science+Business Media, LLC, part of Springer Nature 2018

Correction to: Surgical Endoscopy
<https://doi.org/10.1007/s00464-018-6057-8>

In the original article, there are two errors in Table 3:

1. There were 8 patients undergoing VATE suffer from moderate fever, not 9.
2. In the fourth line of outcomes, saying “pneumothorax, moderate fever and moderate fever,” “moderate fever” was repeated.

The corrected Table 3 is shown here. The authors regret the errors.

Table 3 Safety outcomes in the STER and VATE groups

Outcomes	STER (n = 30)	VATE (n = 28)
Subcutaneous or mediastinal emphysema	3	0
Moderate fever	1	8
Severe chest pain	0	1
Pneumothorax and moderate fever	1	0
Moderate fever, severe chest pain and ventricular fibrillation	0	1
Total AEs, n (%)	5 (16.7%)	10 (35.7%)

STER submucosal tunneling endoscopic resection, VATE video-assisted thoracoscopic enucleation, AE adverse event

The original article can be found online at <https://doi.org/10.1007/s00464-018-6057-8>.

✉ Enqiang Linghu
linghuenqiang@vip.sina.com

¹ Department of Gastroenterology and Hepatology, Chinese PLA General Hospital, Fuxing Road 28. Haidian District, Beijing, People's Republic of China

² Department of Thoracic Surgery, Chinese PLA General Hospital, Beijing, People's Republic of China