ERRATUM

Ahmet Nayir

Circumcision for the prevention of significant bacteriuria in boys

Pediatr Nephrol (2001) 16:1129–1134

In the above article, Table 4 contains erroneous information. “*Enterobacter*” should appear instead of “Enterococci”.

Table 4 Bacterial isolates in the urine cultures (*P. mirabilis* *Proteus mirabilis*, *E. coli* *Escherichia coli*, *S. aureus* *Staphylococcus aureus*, *P. aeruginosa* *Pseudomonas aeruginosa*, *P. vulgaris* *Proteus vulgaris*, *K. oxytoca* *Klebsiella oxytoca*)

	Urinary bacterial isolates in the first urine culture	Urinary bacterial isolates in the first 6-month period	Urinary bacterial isolates in the second 6-month period
Group 1	Total: 35 positive urine cultures <i>P. mirabilis</i> (18) 51.4% <i>E. coli</i> (7) 20% <i>P. aeruginosa</i> (3) 8.5% <i>Enterobacter</i> (2) 5.7% α -Streptococci (2) 5.7% <i>P. vulgaris</i> (1) 5.7% <i>K. oxytoca</i> (1) 5.7% <i>Citrobacter</i> (1) 5.7%	Total: 210 urine cultures positive: 125 urine cultures (59.5%) <i>P. mirabilis</i> (72) 57.6% <i>E. coli</i> (24) 19.2% <i>S. aureus</i> (4) 3.2% <i>P. aeruginosa</i> (5) 4% <i>Enterobacter</i> (6) 4.8% α -Streptococci (7) 5.6% <i>P. vulgaris</i> (5) 4% <i>K. oxytoca</i> (1) 0.8% <i>Citrobacter</i> (1) 0.8%	Total: 210 urine cultures positive: 5 urine cultures (2.3%) <i>S. aureus</i> (4) 80% <i>E. coli</i> (1) 20%
Group 2	Total: 35 positive urine cultures <i>P. mirabilis</i> (20) 57.1% <i>E. coli</i> (6) 17.1% <i>P. aeruginosa</i> (2) 5.7% <i>Enterobacter</i> (2) 5.7% α -Streptococci (3) 8.6% <i>K. oxytoca</i> (2) 5.7%	Total: 210 urine cultures positive: 6 urine cultures (2.8%) <i>S. aureus</i> (3) 50% <i>S. albus</i> (1) 16.6% <i>P. mirabilis</i> (1) 16.6% <i>E. coli</i> (1) 16.6%	—
Uncircumcised group	Total: 18 positive urine cultures <i>P. mirabilis</i> (11) 61% <i>E. coli</i> (4) 22% α -Streptococci (1) 5.5% <i>K. oxytoca</i> (1) 5.5% <i>Enterobacter</i> (1) 5.5%	Total: 108 urine cultures positive: 62 urine cultures (57.4%) <i>P. mirabilis</i> (36) 58% <i>E. coli</i> (12) 19.3% <i>S. aureus</i> (2) 2.7% <i>P. aeruginosa</i> (1) 1.8% <i>Enterobacter</i> (6) 9.6% α -Streptococci (3) 4.8% <i>K. oxytoca</i> (2) 3.2%	Total: 108 urine cultures positive: 60 urine cultures (55.5%) <i>P. mirabilis</i> (35) 58.3% <i>E. coli</i> (14) 23.3% <i>S. aureus</i> (2) 3.3% <i>P. aeruginosa</i> (2) 3.3% <i>Enterobacter</i> (5) 8.3% α -Streptococci (2) 3.3%

A. Nayir

Department of Pediatrics, Pediatric Nephrology Division,
Istanbul Faculty of Medicine, University of Istanbul,
Tarik Zafer Tunaya sok 2/6, Gumussuyu-Istanbul, 80040 Turkey
e-mail: nayir@ttnet.net.tr
Tel.: +90-212-6211644, Fax: +90-212-6211643