Riitta Ojala · Marja Ala-Houhala ·
Aimo P. T. Harmoinen · Tiina Luukkaala ·
Jukka Uotila · Outi Tammela

Tubular proteinuria in pre-term and full-term infants

Published online: 20 July 2006
© IPNA 2006

The online version of the original article can be found at: <http://dx.doi.org/10.1007/s00467-005-2028-0>.

R. Ojala (✉) · M. Ala-Houhala · O. Tammela
Department of Pediatrics, Tampere University Hospital,
P.O. Box 2000, 33521 Tampere, Finland
e-mail: riitta.ojala@pshp.fi
Tel.: +358-3-3116111
Fax: +358-3-2475314

J. Uotila
Department of Gynecology, Tampere University Hospital,
Tampere, Finland

R. Ojala
Pediatric Research Center, Tampere University Hospital,
Tampere, Finland

A. P. T. Harmoinen
Department of Clinical Chemistry,
Savonlinna Central Hospital,
Savonlinna, Finland

T. Luukkaala
School of Public Health Research Unit,
Tampere University Hospital,
Tampere, Finland

Pediatr Nephrol (2006) 21:68–73

There is an error in the first paragraph of the Results section of this manuscript. The correlation between first urinary alpha 1-M/creatinine ratio and gestational age (line 6) should be read -0.647 .

The authors apologize for this error.