Familial steroid-sensitive nephrotic syndrome in Southern Israel: clinical and genetic observations

Daniel Landau · Tal Oved · Dan Geiger · Luba Abizov ·
Hanna Shalev · Ruti Parvari

Published online: 20 February 2007
© IPNA 2007

Unfortunately the column headings were incorrectly titled.
The correct Table 1 is printed below:

Table 1 Patients' characteristics (*SS* simple steroid-sensitive, *FR* frequent-relapsing, *SD* steroid-dependent, *NA* not available, *NS* not significant)

Characteristics	Jews	Bedouins		<i>P</i> value ^a
		Index family		
Number	48	48	11	
Male (%)	37 (78)	38 (77)	8 (73)	NS
Family history (%)	2 (4.2)	13 (28)	NA	<0.005
Age at onset (years)	4.8±3.4	4.9±2.5	4.4±2.4	NS
NS Type				
SS (%)	19 (40)	34 (71)	4 (36)	<0.005
FR (%)	16 (33)	10 (20)	1 (9)	0.08
SD (%)	12 (25)	4 (8)	4 (27)	0.13
NA (%)	1 (2)	0	2 (18)	NS
Steroid-sparing therapy (%)	17 (35)	10 (21)	1 (10)	0.05
Major complications (%)	8 (18)	8 (17)	4 (36)	NS

^a Values refer to comparison between Bs and Js

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00467-006-0409-7>.

D. Landau · H. Shalev
Department of Pediatrics, Faculty of Health Sciences,
Ben Gurion University of the Negev,
Beer-Sheva, Israel

T. Oved · L. Abizov · R. Parvari
Department of Developmental Genetics and Virology,
Faculty of Health Sciences, Ben Gurion University of the Negev,
Beer-Sheva, Israel

D. Geiger
Computer Science Department, Technion,
Haifa, Israel

D. Landau (✉) · H. Shalev
Department of Pediatrics, Soroka Medical Center,
P.O. Box 151, Beer-Sheva 84101, Israel
e-mail: ldaniel@bgu.ac.il