Erratum to: Cyclosporine A vs. methylprednisolone for Henoch–Schönlein nephritis: a randomized trial

Outi Jauhola · Jaana Ronkainen · Helena Autio-Harmainen · Olli Koskimies ·
Marja Ala-Houhala · Pekka Arikoski · Tuula Hölttä · Timo Jahnukainen ·
Jukka Rajantie · Timo Örmälä · Matti Nuutinen

Published online: 4 August 2011
© IPNA 2011

Erratum to: Pediatric Nephrology DOI 10.1007/s00467-011-1919-5

Unfortunately Table 1 was printed incorrectly. The correct version can be seen below.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00467-011-1919-5>.

O. Jauhola · M. Nuutinen (✉)
Department of Children and Adolescents, Oulu University
Hospital,
PO Box 23, 90029, Oulu, Finland
e-mail: matti.nuutinen@ppshp.fi

O. Jauhola
e-mail: ojauhola@paju.oulu.fi

J. Ronkainen
Oulu City Health Care Centre,
Oulu, Finland

H. Autio-Harmainen
Department of Pathology, Oulu University Hospital,
Oulu, Finland

O. Koskimies · T. Hölttä · T. Jahnukainen
Hospital for Children and Adolescents, University of Helsinki,
Helsinki, Finland

M. Ala-Houhala
Department of Pediatrics, Tampere University Hospital,
Tampere, Finland

P. Arikoski
Department of Pediatrics, Kuopio University Hospital,
Kuopio, Finland

J. Rajantie
Department of Pediatrics, Helsinki University Central Hospital,
Jorvi Hospital,
Espoo, Finland

T. Örmälä
Department of Pediatrics, Hyvinkää Hospital,
Hyvinkää, Finland

Table 1 Demographic and clinical features of 24 patients with HSN at onset by treatment group and randomization

	CyA (<i>n</i> =11)		MP (<i>n</i> =13)	
	Randomized (<i>n</i> =7)	Non-randomized (<i>n</i> =4)	Randomized (<i>n</i> =8)	Non-randomized (<i>n</i> =5)
Gender	2 girls, 5 boys	1 girl, 3 boys	2 girls, 6 boys	4 girls, 1 boy
Age at onset (years)				
Mean	9.0	10.4	7.9	11.5
Range	5.1–12.3	7.0–14.4	4.0–14.8	7.8–16.8
dU-Prot (mg/m ² /h)				
Mean	143	73	182	128
Range	38–510	14–229	7–356	83–181
SAlb (g/l)				
Mean	33	31	26	27
Range	20–43	16–40	15–37	14–35
Hematuria N/%	7/7 (100%)	4/4 (100%)	8/8 (100%)	4/4 (100%)
Time from HSP diagnosis to enrollment (months)				
Mean	3.0	9.2	3.6	4.6
Range	0.8–11.8	1.7–30.6	0.9–7.8	1.5–8.5
ISKDC II	2	0	3	0
ISKDC III	5	4	5	5