Mark G. Kris
Paul J. Hesketh
Jorn Herrstedt
Cynthia Rittenberg
Lawrence H. Einhorn
Steven Grunberg
Jim Koeller
Ian Olver
Sussanne Borjeson
Enzo Ballatori

Consensus proposals for the prevention of acute and delayed vomiting and nausea following high-emetic-risk chemotherapy

Published online: 10 May 2005
© Springer-Verlag 2005

Support Care Cancer (2005) 13:85–96

In Table 1 the dosage of dolasetron was incorrectly given as 0.18 mg/kg. The correct dosage is 1.8 mg/kg.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00520-004-0699-x>

M. G. Kris (✉)
Memorial Sloan-Kettering Cancer Center,
1275 York Avenue,
New York, NY, 10021, USA
e-mail: krism@mskcc.org
Tel.: +1-212-6397590
Fax: +1-212-7944357

P. J. Hesketh
Tufts University School of Medicine,
Boston, MA, USA

J. Herrstedt
Copenhagen University Hospital,
Copenhagen, Denmark

C. Rittenberg
Rittenberg Oncology Consulting,
Metairie, LA, USA

L. H. Einhorn
Walther Cancer Institute,
Indianapolis, IN, USA

S. Grunberg
Vermont Cancer Center,
Shelburne, VT, USA

J. Koeller
University of Texas,
Austin, TX, USA

I. Olver
Royal Adelaide Hospital Cancer Center,
Adelaide, SA, Australia

S. Borjeson
Linköping University,
Linköping, Sweden

E. Ballatori
Oncologia Medica,
Perugia, Italy