A meta-analysis of exercise interventions among people treated for cancer

Vicki S. Conn • Adam R. Hafdahl • Davina C. Porock •
Roxanne McDaniel • Paul J. Nielsen

Published online: 8 September 2007
© Springer-Verlag 2007

Erratum to: Support Care Cancer DOI 10.1007/s00520-005-0905-5

In the article “A Meta-Analysis of Exercise Interventions among People Treated for Cancer,” incorrect quality of life results were reported due to a coding error of baseline values for one primary study. The error affected quality of life outcome data for the treatment group baseline versus post intervention comparison and for the control group baseline

versus outcome comparison. The corrected data appear in the tables below.

One article sentence should be corrected (p. 705): The original sentence was “All ESs are small and statistically nonsignificant, except for quality of life at -0.26 under the assumption of no pre- and postscore association”; it should read “All ESs are small and statistically nonsignificant.” The authors apologize for the error.

Tables 4 and 5

The online version of this article can be found at <http://dx.doi.org/10.1007/s00520-005-0905-5>.

V. S. Conn (✉)
S317 School of Nursing, University of Missouri–Columbia,
Columbia, MO 65211, USA
e-mail: conn@missouri.edu

A. R. Hafdahl
28 McAlester Hall, Department of Psychological Sciences,
University of Missouri–Columbia,
Columbia, MO 65211, USA

D. C. Porock
Queens Medical Centre, Nottingham Education Center,
University of Nottingham,
NG7 2HA Nottingham, UK

R. McDaniel
S410 School of Nursing, University of Missouri–Columbia,
Columbia, MO 65211, USA

P. J. Nielsen
Biochemistry Department, University of Missouri–Columbia,
Columbia, MO 65211, USA

Table 4 Fixed- and random-effects results by outcome measure: single-group comparisons of pretest and post on quality of life

Association assumed	k	Fixed-effects analysis				Random-effects analysis			
		$\bar{d}$	$SE(\bar{d})$	95% CI for δ	Q	$\hat{\mu}_{\delta}$	$SE(\hat{\mu}_{\delta})$	95% CI for μ_{δ}	$\hat{\sigma}_{\delta}$
$\rho_{12} = .00$	14	0.37***	0.074	0.22–0.51	8.58	-	-	-	0.000 ^a
$\rho_{12} = .80$	15	0.38***	0.034	0.31–0.45	49.09***	0.45***	0.068	0.31–0.58	0.214

Table 5 Fixed- and random-effects results by outcome measure: single-group pre-post comparisons for control subjects on quality of life

Association assumed	k	Fixed-effects analysis				Random-effects analysis			
		$\bar{d}$	$SE(\bar{d})$	95% CI for δ	Q	$\hat{\mu}_{\delta}$	$SE(\hat{\mu}_{\delta})$	95% CI for μ_{δ}	$\hat{\sigma}_{\delta}$
$\rho_{12} = .00$	5	-0.09	0.112	-0.31–0.13	1.03	-	-	-	0.000 ^a
$\rho_{12} = .80$	5	-0.09 [†]	0.050	-0.19–0.01	5.09	-0.08	0.061	-0.20–0.04	0.063