Aqua lymphatic therapy in women who suffer from breast cancer treatment-related lymphedema: a randomized controlled study

Dorit Tidhar · Michal Katz-Leurer

Published online: 17 July 2009
© Springer-Verlag 2009

Erratum to: Support Care Cancer DOI 10.1007/s00520-009-0669-4

Page 5 second paragraph:

“Quality of life was assessed by a questionnaire that was developed by **Mayrovitz et al.**”

On page 5 paragraph 3:

“The questionnaire was found to be reliable and valid for breast cancer lymphedema patients. **Mayrovitz et al.**”

On page 8 at the end of paragraph 2:

“and moderate according to **Mayrovitz et al.** [27].”

All three references should be replaced by: **Launois et al.**

An incorrect version of Table 3 was published. The correct table is given here.

Table 3 Baseline and final volume of the lymphedema arm in the first and last ALT sessions in the study group

	Volume (ml)		<i>p</i> value ^a
	Presession	Postsession	
First session	2,513.5 (695.7)	2,459.2 (693.1)	0.02
Last session	2,613.9 (846.70)	2,515.2 (865.8)	<0.01

Data are reported as mean (SD)

^aAs determined by pair *t* test

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00520-009-0669-4>

D. Tidhar

Department of Physical Therapy, “Maccabi Healthcare Services”,
Netivot, Israel

M. Katz-Leurer (✉)

Department of Physical Therapy, School of Health Professions,
Sackler Faculty of Medicine, Tel-Aviv University,
Tel-Aviv 69978, Israel
e-mail: michalkz@post.tau.ac.il