Erratum to: Effects of the enteral administration of *Bifidobacterium breve* on patients undergoing chemotherapy for pediatric malignancies

Mariko Wada · Satoru Nagata · Masahiro Saito ·
Toshiaki Shimizu · Yuichiro Yamashiro ·
Takahiro Matsuki · Takashi Asahara · Koji Nomoto

Published online: 26 June 2010
© Springer-Verlag 2010

Erratum to: Support Care Cancer
DOI 10.1007/s00520-009-0711-6

Unfortunately, an error occurred in Table 3 of the article.
Please find below the corrected table.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00520-009-0711-6>.

M. Wada · S. Nagata · Y. Yamashiro · K. Nomoto
Department of Laboratory for Probiotics Research,
Juntendo University School of Medicine,
Tokyo 113-8421, Japan

M. Wada (✉) · S. Nagata · M. Saito · T. Shimizu
Department of Pediatrics,
Juntendo University School of Medicine,
1-1, Hongo 2-chome, Bunkyo-ku,
Tokyo 113-8421, Japan
e-mail: maro@juntendo.ac.jp

T. Matsuki · T. Asahara · K. Nomoto
Yakult Central Institute for Microbiological Research,
Tokyo 186-8650, Japan

Table 3 Fecal microflora after the administration of the study products

Weeks after the beginning of intervention												
	0W		2W		3W		4W		5W		6W	
	Placebo	<i>B. breve</i>	Placebo	<i>B. breve</i>	Placebo	<i>B. breve</i>	Placebo	<i>B. breve</i>	Placebo	<i>B. breve</i>	Placebo	<i>B. breve</i>
Anaerobes												
Total anaerobes	9.8±0.7 (4/4)	10.1±0.8 (11/11)	9.6±0.6 (7/7)	10.2±0.4 (9/9)	9.7±0.3 (8/8)	10.1±0.5 (7/7)	9.9±0.5 (13/13)	9.9±0.9 (13/13)	9.6±0.9 (7/7)	9.9±0.9 (9/9)	9.8±0.5 (7/9)	10.5±0.1 (2/2)
Clostridium leptum subgroup	8.9±0.7 (4/5)	9.2±1.1 (13/13)	8.1±0.8 (9/9)	9.4±0.5* (9/9)	8.8±0.6 (9/10)	8.1±1.3 (9/10)	9.0±0.7 (13/14)	9.0±0.9 (12/13)	8.7±1.4 (9/10)	9.0±1.3 (7/9)	8.7±0.9 (9/11)	9.5±0.5 (6/6)
<i>B. stain</i> Yakult	ND (0/4)	ND (0/11)	ND (0/7)	6.1±2.3 c (6/9)	ND (0/10)	6.6±1.0 c (7/7)	ND (0/13)	5.9±1.3 c (12/13)	ND (0/7)	6.8±2.0 c (8/8)	ND (0/9)	8.0 c (1/2)
Facultative anaerobes												
Total facultative anaerobes	8.4±0.9 (4/4)	8.5±0.8 (11/11)	9.2±0.8 (7/7)	8.0±1.0 (9/9)	8.8±1.1 (10/10)	9.1±0.8 (7/7)	8.8±1.0 (13/13)	8.5±0.8 (13/13)	8.6±1.3 (7/7)	8.4±0.9 (9/9)	8.7±1.0 (9/9)	8.3 (2/2)
Enterobacteriaceae	5.6±0.8 (2/4)	6.1±2.5 (10/11)	6.6±1.2 (7/7)	5.7±2.8 (5/7)	7.9±1.7 b (8/10)	7.5±0.9 (6/7)	6.6±2.4 b (10/13)	6.6±2.0 (8/13)	5.8±2.7 (4/7)	7.0±1.9 (7/9)	6.5±2.3 (6/9)	9.0 (1/2)
Enterococcus	7.4±1.3 (4/4)	8.2±1.5 (11/11)	9.1±1.0 (7/7)	7.5±0.9 a (9/9)	8.6±1.1 (10/10)	8.7±1.2 (7/7)	8.6±1.3 (13/13)	8.2±1.1 (13/13)	8.2±1.6 (7/7)	8.2±1.2 (9/9)	8.7±1.7 (9/9)	8.2 (2/2)
Aerobes												
<i>Bacillus</i>	5.7±2.0 (3/4)	5.1±1.7 (10/11)	3.2 (1/7)	4.9±1.6 (7/9)	5.3±2.4 (7/10)	5.1±2.0 (5/7)	4.4±1.4 (8/13)	5.8±2.2 (8/13)	5.3±2.0 (5/7)	4.5±1.7 b (7/9)	5.1±1.7 (7/9)	2.9 (2/2)
<i>Pseudomonas</i>	ND (0/4)	ND (0/11)	ND (0/7)	ND (0/9)	ND (0/10)	ND (1/7)	ND (0/13)	4.0 (1/13)	ND (0/7)	ND (0/9)	ND (0/9)	ND (0/2)

Values are mean ± SD (Real-time PCR, log₁₀ cells/g of feces; Culture technique, log₁₀ CFU/g of feces)

ND, Not detected (Real-time PCR, <6.0 log₁₀ cells/g of feces; Culture technique, <1.6 - <2.9 log₁₀ CFU/g of feces)

(), Incidence (number of patients detected/number of patients measured)

Statistical Software: SPSS Ver. 11 (SPSS Japan Inc.), 2-sided *p*-value<0.05

Unpaired-t test:**p*<0.05, ** *p*<0.01

Paired-t test: 0 W vs 2 W;a *p*<0.05, 2Wvs 3,4,5 or 6 W; b *p*<0.05, bb *p*<0.01

Fisher's exact test: c *p*<0.01