Erratum to: Hepatitis B virus management to prevent reactivation after chemotherapy: a review

Jessica P. Hwang · John M. Vierling ·
Andrew D. Zelenetz · Susan C. Lackey · Rohit Loomba

Published online: 27 January 2013
© Springer-Verlag Berlin Heidelberg 2013

**Erratum to: Support Care Cancer (2012) 20:2999-3008
DOI 10.1007/s00520-012-1576-7**

The original version of this article unfortunately contained mistake. There was an error in citing reference [54]. The paragraph reads “Other chemotherapies used to treat hema-

tologic malignancies that have been associated with reactivation include CHOP (cyclophosphamide, doxorubicin, vincristine, and prednisone) for NHL [52], bendamustine for chronic lymphocytic leukemia (CLL) [53] and NHL [54], chlorambucil for CLL [55], and imatinib mesylate for myeloid leukemia [56].” NHL [54] should be replaced by multiple myeloma [54].

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00520-012-1576-7>.

J. P. Hwang · S. C. Lackey
The University of Texas MD Anderson Cancer Center,
Houston, TX, USA

J. M. Vierling
Baylor College of Medicine,
Houston, TX, USA

A. D. Zelenetz
Memorial Sloan-Kettering Cancer Center,
New York City, NY, USA

R. Loomba
The University of California San Diego,
La Jolla, CA, USA

J. P. Hwang (✉)
Department of General Internal Medicine,
1515 Holcombe Blvd., Unit 1465,
Houston, TX 77030, USA
e-mail: jphwang@mdanderson.org