Erratum to: Risk factors for recurrence after transarterial chemoembolization for early-stage hepatocellular carcinoma

Hideaki Kinugasa · Kazuhiro Nouse · Yasuto Takeuchi · Tetsuya Yasunaka · Hideki Onishi · Shin-ichiro Nakamura · Hidenori Shiraha · Kenji Kuwaki · Hiroaki Hagihara · Fusao Ikeda · Yasuhiro Miyake · Akinobu Takaki · Kazuhide Yamamoto

Published online: 15 December 2011
© Springer 2011

Erratum to: J Gastroenterol DOI 10.1007/s00535-011-0492-9

Please note that errors appeared in the above-cited article.

In Abstract, in the “Results” section the first sentence should read: “The local recurrence rates and intrahepatic distant recurrence rates at 3 months, 6 months, and 1 year were 18.6, 33.4, and 61.8%, and 2.8, 2.8, and 34.3%, respectively.”

In the main text, in the Results section, under the heading “Recurrence rate”, in the second paragraph the

second sentence should read: “The local recurrence rates and intrahepatic distant recurrence rates at 3 months, 6 months, and 1 year were 18.6, 33.4, and 61.8%, and 2.8, 2.8, and 34.3%, respectively (Fig. 1).”

In the legend for Fig. 1, the second sentence should read: “The local recurrence rates (*solid line*) and intrahepatic distant recurrence rates (*dotted line*) at 3 months, 6 months, and 1 year were 18.6, 33.4, and 61.8%, and 2.8, 2.8, and 34.3%, respectively.”

The online version of the original article can be found under doi:[10.1007/s00535-011-0492-9](https://doi.org/10.1007/s00535-011-0492-9).

H. Kinugasa (✉) · K. Nouse · Y. Takeuchi · T. Yasunaka · H. Onishi · S. Nakamura · H. Shiraha · K. Kuwaki · H. Hagihara · F. Ikeda · Y. Miyake · A. Takaki · K. Yamamoto
Department of Gastroenterology and Hepatology,
Okayama University Graduate School of Medicine, Dentistry,
and Pharmaceutical Sciences, 2-5-1 Shikata-cho,
Kita-ku, Okayama 700-8558, Japan
e-mail: gyacy14@gmail.com