CORRECTION

Correction to: Complete mucosal healing of distal lesions induced by twice-daily budesonide 2-mg foam promoted clinical remission of mild-to-moderate ulcerative colitis with distal active inflammation: double-blind, randomized study

Makoto Naganuma¹ · Nobuo Aoyama² · Tomohiro Tada³ · Kiyonori Kobayashi⁴ · Fumihito Hirai⁵ · Kenji Watanabe⁶ · Mamoru Watanabe⁷ · Toshifumi Hibi⁸

Published online: 13 October 2017

© Japanese Society of Gastroenterology 2017

Correction to: J Gastroenterol

DOI 10.1007/s00535-017-1376-4

In Fig. 2(c) of this article, the length of the “95% CI difference bars” under the items labelled “Left-sided and Pancolitis” and “Left-sided” under the heading “Disease type” was corrected. The correct figure is shown in the following page.

The online version of the original article can be found under doi:[10.1007/s00535-017-1376-4](https://doi.org/10.1007/s00535-017-1376-4).

✉ Makoto Naganuma
maknaganuma@gmail.com

- ¹ Department of Gastroenterology and Hepatology, School of Medicine, Keio University, Shinanomachi 35, Shinjuku-ku, Tokyo 160-0016, Japan
- ² Gastrointestinal Endoscopy and Inflammatory Bowel Disease Center, Aoyama Medical Clinic, Hyogo, Japan
- ³ Tada Tomohiro Institute of Gastroenterology and Proctology, Saitama, Japan
- ⁴ Department of Research and Development Center for New Medical Frontiers, School of Medicine, Kitasato University, Kanagawa, Japan
- ⁵ Department of Gastroenterology, Fukuoka University Chikushi Hospital, Fukuoka, Japan
- ⁶ Department of Gastroenterology, Osaka City General Hospital, Osaka, Japan
- ⁷ Department of Gastroenterology and Hepatology, School of Medicine, Tokyo Medical and Dental University, Tokyo, Japan
- ⁸ Center for Advanced IBD Research and Treatment, Kitasato Institute Hospital, Kitasato University, Tokyo, Japan

(d)

		Clinical remission, n/N (%)				
		Overall	Proctitis	Left-sided/Pancolitis	Left-sided	Pancolitis
Complete mucosal healing	Achieved	16/21 (76.2)	5/7 (71.4)	11/14 (78.6)	9/11 (81.8)	2/3 (66.7)
	Not achieved	11/43 (25.6)	5/15 (33.3)	6/28 (21.4)	4/20 (20.0)	6/28 (21.4)

In Fig. 3(a) of this article, the length of the “95% CI difference bar” for the items labelled “Yes” under the heading “Rectal 5-ASA” was corrected. In Fig. 3(b), the length of the “95% CI difference bars” for the items labelled “Yes” under the heading “Rectal 5-ASA”, the

item “Low” under the heading “Dose of oral 5-ASA”, and the item “6≤” under the heading “Baseline MMDAI” was corrected. The correct figure is shown in the following page.