Sickness absence and concurrent low back and neck–shoulder pain: results from the MUSIC-Norrtälje study

Teresia Nyman · Wilhelmus Johannes Andreas Grooten ·
Christina Wiktorin · Johan Liwing · Linda Norrman

Published online: 17 November 2006
© Springer-Verlag 2006

Erratum to: Eur Spine J DOI 10.1007/s00586-006-0152-6

Due to an error on the part of the typesetter, the numbering of the reference citations in the text of this article did not correspond to the numbering of the reference list. The correctly numbered reference list is given here.

References

1. Allebeck P, Mestekaasa A (2004) Swedish Council on Technology Assessment in Health Care (SBU). Chapter 3. Causes of sickness absence: research approaches and explanatory models. *Scand J Public Health Suppl* 63:36–43
2. Andersson GB (1999) Epidemiological features of chronic low-back pain. *Lancet* 354:581–585
3. Borg K, Hensing G, Alexanderson K (2001) Predictive factors for disability pension—an 11-year follow up of young persons on sick leave due to neck, shoulder, or back diagnoses. *Scand J Public Health* 29:104–112
4. Borghouts JA, Koes BW, Vondeling H et al (1999) Cost-of-illness of neck pain in The Netherlands in 1996. *Pain* 80:629–636
5. Cote P, Cassidy JD, Carroll LJ et al (2004) The annual incidence and course of neck pain in the general population: a population-based cohort study. *Pain* 112:267–273
6. Florwall Muller CF, Monrad T, Biering-Sorensen F et al (1999) The influence of previous low back trouble, general health, and working conditions on future sick-listing because of low back trouble. A 15-year follow-up study of risk indicators for self-reported sick-listing caused by low back trouble. *Spine* 24:1562–1570
7. Fransen M, Woodward M, Norton R et al (2002) Risk factors associated with the transition from acute to chronic occupational back pain. *Spine* 27:92–98
8. Goldberg D, Williams P (1988) A user's guide to the general health questionnaire. NFER-NELSON, Windsor
9. Grooten WJ, Wiktorin C, Norrman L et al (2004) Seeking care for neck/shoulder pain: a prospective study of work-related risk factors in a healthy population. *J Occup Environ Med* 46:138–146
10. Hansson EK, Hansson TH (2005) The costs for persons sick-listed more than one month because of low back or neck problems. A two-year prospective study of Swedish patients. *Eur Spine J* 14:337–345
11. Hansson T, Jensen I (2004) Swedish Council on Technology Assessment in Health Care (SBU). Chapter 6. Sickness absence due to back and neck

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00586-006-0152-6>.

T. Nyman (✉) · W. J. A. Grooten · C. Wiktorin ·
J. Liwing · L. Norrman
Department of Occupational and Environmental Health,
Stockholm County Council, Norrbacka,
171 76 Stockholm, Sweden
e-mail: teresia.nyman@sl.se

T. Nyman · W. J. A. Grooten · C. Wiktorin
Department of Public Health Sciences,
Division of Occupational Medicine,
Karolinska Institutet, Stockholm, Sweden

- disorders. *Scand J Public Health Suppl* 63:109–151
12. Hensing G (2004) Swedish Council on Technology Assessment in Health Care (SBU). Chapter 4. Methodological aspects in sickness-absence research. *Scand J Public Health Suppl* 63:44–48
13. Hurwitz EL, Morgenstern H (1999) The effect of comorbidity on care seeking for back problems in the United States. *Ann Epidemiol* 9:262–270
14. IJzelenberg W, Burdorf A (2004) Impact of musculoskeletal co-morbidity of neck and upper extremities on healthcare utilisation and sickness absence for low back pain. *Occup Environ Med* 61:806–810
15. Kecklund G, Åkerstedt T (1992) The psychometric properties of the Karolinska Sleep Questionnaire. *J Sleep Res* 1(Suppl):133
16. Nachemson A, Waddell G, Norlund A (2000) Epidemiology of neck and low back pain. In: Nachemson A, Jonsson E (eds) *Neck and back pain: the scientific evidence of causes, diagnosis and treatment*. Lippincott, Williams & Wilkins, Philadelphia, pp 165–187
17. National Social Insurance Board (2003) Vad kostar sjukskrivningarna inom olika yrken? RFV redovisar 2003:5 [Swedish]. National Social Insurance Board, Stockholm
18. National Social Insurance Board (2004) Vad kostar sjukdomarna för kvinnor och män? RFV redovisar 2004:5 [Swedish]. National Social Insurance Board, Stockholm
19. Nordin M, Hiebert R, Pietrek M et al (2002) Association of comorbidity and outcome in episodes of nonspecific low back pain in occupational populations. *J Occup Environ Med* 44:677–684
20. Pernold G, Mortimer M, Wiktorin C et al (2005) Neck/shoulder disorders in a general population. Natural course and influence of physical exercise: a 5-year follow-up. *Spine* 30:E363–E368
21. Picavet HS, Schouten JS (2003) Musculoskeletal pain in the Netherlands: prevalences, consequences and risk groups, the DMC(3)-study. *Pain* 102:167–178
22. Seferlis T, Nemeth G, Carlsson AM et al (1999) Acute low-back-pain patients exhibit a fourfold increase in sick leave for other disorders: a case-control study. *J Spinal Disord* 12:280–286
23. Severens JL, Mulder J, Laheij RJ et al (2000) Precision and accuracy in measuring absence from work as a basis for calculating productivity costs in The Netherlands. *Soc Sci Med* 52:243–249
24. Theorell T, Ahlberg-Hulten G, Sigala F et al (1990) A psychosocial and biomedical comparison between men in six contrasting service occupations. *Work Stress* 4:51–63
25. Theorell T, Konarski-Svensson JK, Ahlmen J et al (1991) The role of paid work in Swedish chronic dialysis patients—a nation-wide survey: paid work and dialysis. *J Intern Med* 230:501–509
26. Tornqvist EW, Kilbom A, Vingard E et al (2001) The influence on seeking care because of neck and shoulder disorders from work-related exposures. *Epidemiology* 12:537–545
27. Waddell G (1998) *The back pain revolution*. Churchill Livingstone, London
28. Waldenstrom M, Theorell T, Ahlberg G et al (2002) Assessment of psychological and social current working conditions in epidemiological studies: experiences from the MUSIC-Norrtaälje study. *Scand J Public Health* 30:94–102
29. van der Giezen AM, Bouter LM, Nijhuis FJ (2000) Prediction of return-to-work of low back pain patients sicklisted for 3–4 months. *Pain* 87:285–294
30. van Poppel MN, de Vet HC, Koes BW et al (2002) Measuring sick leave: a comparison of self-reported data on sick leave and data from company records. *Occup Med (Lond)* 52:485–490
31. van Tulder MW, Koes BW, Bouter LM (1995) A cost-of-illness study of back pain in The Netherlands. *Pain* 62:233–240
32. Vingard E, Alexanderson K, Norlund A (2004) Swedish Council on Technology Assessment in Health CARE (SBU). Chapter 9. Consequences of being on sick leave. *Scand J Public Health Suppl* 63:207–215
33. Vingard E, Alfredsson L, Hagberg M et al (2000) To what extent do current and past physical and psychosocial occupational factors explain care-seeking for low back pain in a working population? Results from the Musculoskeletal Intervention Center-Norrtaälje Study. *Spine* 25:493–500
34. Von Korf M, Ormel J, Keefe FJ et al (1992) Grading the severity of chronic pain. *Pain* 50:133–149