The change in ratio of convex and concave lung volume in adolescent idiopathic scoliosis: a 3D CT scan based cross sectional study of effect of severity of curve on convex and concave lung volumes in 99 cases

**Eun Mi Chun · Seung Woo Suh · Hitesh N. Modi ·
Eun Young Kang · Suk Joo Hong · Hae-Ryong Song**

Published online: 18 January 2008
© Springer-Verlag 2008

Erratum to: Eur Spine J
DOI 10.1007/s00586-007-0488-6

The given name of the fifth author was incorrect. The correct spelling is Suk Joo Hong.

An incorrect address for E.Y. Kang and S.J. Hong appeared in the original article. The revised version is given below:

E.Y. Kang · S.J. Hong
Department of Radiology, Korea University,
Guro Hospital, Seoul, South Korea

The online version of the original article can be found under doi:[10.1007/s00586-007-0488-6](https://doi.org/10.1007/s00586-007-0488-6).

S. W. Suh (✉) · H. N. Modi
Department of Orthopaedic Surgery, Korea University,
Guro Hospital, #80, Guro-Dong, Guro-Gu,
Seoul 152-703, South Korea
e-mail: spine@korea.ac.kr

E. M. Chun
Internal Medicine, Ewha Womans University,
Seoul, South Korea

H.-R. Song
Department of Orthopedics, Korea University,
Guro Hospital, Seoul, South Korea

E. Y. Kang · S. J. Hong
Department of Radiology, Korea University,
Guro Hospital, Seoul, South Korea