Correction to: The quadriceps active ratio: a dynamic MRI-based assessment of patellar height

Sergio Barroso Rosa^{1,2} · Zaid Bahho¹ · Kenji Doma¹ · Kaushik Hazratwala¹ · Peter McEwen¹ · Varaguna Manoharan¹ · Brent Matthews¹ · Matthew Wilkinson¹

Published online: 31 March 2018

© Springer-Verlag France SAS, part of Springer Nature 2018

Correction to:

European Journal of Orthopaedic Surgery & Traumatology

<https://doi.org/10.1007/s00590-018-2170-6>

In the online version of the original article, one of the white lines was not accurately drawn in Fig. 1.

The revised Fig. 1 is placed here:

The original article can be found online at <https://doi.org/10.1007/s00590-018-2170-6>.

✉ Sergio Barroso Rosa
dr.sbarroso@gmail.com

Zaid Bahho
zaid_bahho@hotmail.com

Kenji Doma
kenji.doma@jcu.edu.au

Kaushik Hazratwala
koshman1@bigpond.com

Peter McEwen
peter@kneesurgeon.com.au

Varaguna Manoharan
vmanoharan86@gmail.com

Brent Matthews
drbrentmatthews@gmail.com

Matthew Wilkinson
mprwilkinson@hotmail.com

¹ The ORIQL (Orthopaedic Research Institute of Queensland),
7 Tuner Street, Pimlico, QLD 4812, Australia

² Univeristy of Las Palmas de Gran Canaria, Las Palmas,
Canary Islands, Spain

Fig. 1 Calculation of the PTI—patellotrochlear index (quads active ratio if assessed under active contraction of the quadriceps muscles)