Normal fasting plasma glucose levels and type 2 diabetes: the high-risk and population strategy for occupational health promotion (HIPOP-OHP) study

**Y. Hayashino · S. Fukuhara · Y. Suzukamo ·
T. Okamura · T. Tanaka · H. Ueshima**

Published online: 2 October 2007
© Springer-Verlag 2007

**Erratum to: Acta Diabetol (2007) 44:164–166
DOI 10.1007/s00592-007-0258-2**

The original version of this article unfortunately contained a mistake. The title of the article was incorrectly given. The correct title is as given below.

Normal fasting plasma glucose levels and type 2 diabetes: the high-risk and population strategy for occupational health promotion (HIPOP-OHP) study

The online version of the original article can be found under doi:[10.1007/s00592-007-0258-2](https://doi.org/10.1007/s00592-007-0258-2).

Y. Hayashino (✉) · S. Fukuhara
Department of Epidemiology and Healthcare Research,
Kyoto University Graduate School of Medicine,
Konoe-cho, Yoshida, Sakyo-ku, Kyoto 606-8501, Japan
e-mail: Hayasino-y@umin.net

Y. Suzukamo
Department of Physical Medicine and Rehabilitation,
Tohoku University Graduate School of Medicine,
Sendai, Japan

T. Okamura · T. Tanaka · H. Ueshima
Department of Health Science,
Shiga University of Medical Science, Shiga, Japan