TNF-alpha-238, -308 polymorphisms, and diabetes

Rennan Feng · Ying Li · Changhao Sun

Received: 27 April 2011 / Accepted: 30 April 2011 / Published online: 11 May 2011
© Springer-Verlag 2011

Dear Editor,

Three recent meta-analyses regarding TNF-alpha-238, -308 polymorphisms, and diabetes have been published, which reported that TNF-alpha-238, -308 polymorphisms were not associated with type 2 diabetes mellitus (T2DM); however, TNF-alpha-308 polymorphism was positively associated with type 1 diabetes (T1DM) [1–3]. These findings were interesting but complicated. Several points should be further discussed on these findings.

First, it has been still unclear whether these two single nucleotide polymorphisms (SNPs) could lead to a higher or lower TNF-alpha production in the human beings, although the in vitro experiments results reported that these SNPs regulated the TNF-alpha expression. Seldom population study reported the differences of TNF-alpha between the genotypes of the SNPs. It is important for readers to know whether these SNPs could regulate TNF-alpha secretion in human beings. The determination of TNF-alpha in patients with diabetes with various genotypes of these SNPs could provide readers more information about their roles in the development of diabetes.

Second, a positive association between TNF-alpha and T1DM suggested us that this SNP might exert its role by interacting with other genetic factors in the T1DM development. Previous studies have reported strong linkage disequilibriums between HLA DR3, HLA-DRB1*0301 alleles, and TNF-alpha-308 polymorphism [1, 4], indicating that this significant association of TNF-alpha-308

polymorphism may not be independent. TNF-alpha-308 polymorphism might be only a participant in the whole genetic pathopoiesis of T1DM. Larger range of loci should be considered regarding the interaction between them.

Third, another meta-analysis reported that TNF-alpha-308 polymorphism was associated with metabolic syndrome, and a large-scale study also suggested that smaller effect of this SNP might exist [5, 6], which suggested that TNF-alpha-308 polymorphism might play a role in the development of metabolic disorder; however, its mild effect could not cause T2DM.

References

1. Feng RN, Li Y, Sun CH (2009) TNF 308 G/A polymorphism and type 1 diabetes: a meta-analysis. *Diab Res Clin Pract* 85:e4–e7
2. Feng R, Li Y, Zhao D, Wang C, Niu Y, Sun C (2009) Lack of association between TNF 238 G/A polymorphism and type 2 diabetes: a meta-analysis. *Acta Diabetol* 46:339–343
3. Feng RN, Zhao C, Sun CH, Li Y (2011) Meta-Analysis of TNF 308 G/A Polymorphism and Type 2 Diabetes Mellitus. *PLoS One* 6:e18480
4. Javor J, Ferencik S, Bucova M, Stuchlikova M, Martinka E, Barak L et al (2010) Polymorphisms in the genes encoding TGF-beta1, TNF-alpha, and IL-6 show association with type 1 diabetes mellitus in the Slovak population. *Arch Immunol Ther Exp (Warsz)* 58:385–393
5. Boraska V, Rayner NW, Groves CJ, Frayling TM, Diakite M, Rockett KA et al (2010) Large-scale association analysis of TNF/LTA gene region polymorphisms in type 2 diabetes. *BMC Med Genet* 11:69
6. Sookoian SC, Gonzalez C, Pirola CJ (2005) Meta-analysis on the G-308A tumor necrosis factor alpha gene variant and phenotypes associated with the metabolic syndrome. *Obes Res* 13:2122–2131

R. Feng · Y. Li · C. Sun (✉)

Department of Nutrition and Food Hygiene,
School of Public Health, Harbin Medical University,
157 Baojian Road, Nangang District, Harbin 150086,
People's Republic of China
e-mail: changhao2002sun@gmail.com