Erratum to: Miglitol administered before breakfast increased plasma active glucagon-like peptide-1 (GLP-1) levels after lunch in patients with type 2 diabetes treated with sitagliptin

Kazutaka Aoki · Hiroshi Kamiyama ·
Kouichiro Yoshimura · Makoto Shibuya ·
Kiyomi Masuda · Yasuo Terauchi

Published online: 1 November 2011
© Springer-Verlag 2011

Erratum to: Acta Diabetol DOI 10.1007/s00592-011-0322-9

The original publication of this article contains some typographical errors in the legend of Fig. 5.

The correct version of the legend for Fig. 5 is given below:

Fig. 5 Plasma total GIP levels in miglitol-untreated and miglitol-treated groups. **a** Time profiles of the plasma total GIP levels. **b** AUCs of the plasma total GIP levels from after breakfast until 120 min after lunch for each group. **c** AUCs of the plasma total GIP levels after lunch for each group. The time data are presented as the mean $\pm$ SE. $*P < 0.05$, $**P < 0.01$ and $***P < 0.001$ versus miglitol-untreated group.

The online version of the original article can be found under doi:[10.1007/s00592-011-0322-9](https://doi.org/10.1007/s00592-011-0322-9).

K. Aoki · H. Kamiyama · M. Shibuya · K. Masuda ·
Y. Terauchi (✉)
Department of Endocrinology and Metabolism,
Graduate School of Medicine, Yokohama City University,
3-9 Fukuura, Kanazawa-ku, Yokohama 236-0004, Japan
e-mail: terauchi@yokohama-cu.ac.jp

K. Aoki · H. Kamiyama · K. Yoshimura · M. Shibuya
Department of Endocrinology and Metabolism,
Yokosuka Kyousai Hospital, Yokosuka, Japan