Erratum to: Surgical unroofing of the optic canal and visual outcome in basal meningiomas

Giuseppe Mariniello · Oreste de Divitiis ·
Giulio Bonavolontà · Francesco Maiuri

Published online: 7 October 2012
© Springer-Verlag Wien 2012

Erratum to: Acta Neurochir
DOI 10.1007/s00701-012-1485-z

The original version of this article unfortunately contained mistake. Table 2 was incorrectly presented, correct table 2 is presented below:

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00701-012-1485-z>.

G. Mariniello · O. de Divitiis · F. Maiuri
Department of Neurological Sciences, Section of Neurosurgery,
“Federico II” University School of Medicine,
Naples, Italy

G. Bonavolontà
Department of Ophthalmology, Section of Orbital Pathology,
“Federico II” University School of Medicine,
Naples, Italy

G. Mariniello (✉)
Università degli Studi di Napoli “Federico II”,
Cattedra di Neurochirurgia, 5 Via S. Pansini,
Naples 80131, Italy
e-mail: giumarin@unina.it

Table 2 Summary of the data of 46 clinoidal meningiomas

Findings		Overall (46 cases)	Group A (32 cases)		Group B (14 cases)	
AI-Mefty type	type I	11 (24 %)	8 (25 %)		3 (21.5 %)	
	type II	28 (60.8 %)	20 (62.5 %)		8 (57 %)	
	type III	7 (15.2 %)	4 (12.5 %)		3 (21.5 %)	
tumor size	<2.5 cm	14 (30.4 %)	9 (28.1 %)		5 (35.7 %)	
	2.5 to 4 cm	18 (39.2 %)	12 (37.5 %)		6 (42.9 %)	
	>4 cm	14 (30.4 %)	11 (34.4 %)		3 (21.4 %)	
tumor extension into the OC		10 (22 %)	6 (18.7 %)		4 (28.5 %)	
early ON visualization			<2.5 cm	7/9	<2.5 cm	all
			2.5 to 4 cm	2/12	2.5 to 4 cm	all
			>4 cm	none	>4 cm	all
early ICA visualization			5 (15.6%) (<2.5 cm; types II and III)		14 (100 %)	
complete tumor resection		39 (84.8 %)	26 (81 %)		13 (93 %)	
preoperative visual symptoms		30 (65.2 %)	20 (62.5 %)		10 (71.4 %)	
visual outcome	improved	17 (56.7 %)	9 (45 %)		8 (80 %)	
	stable	12 (40 %)	10 (50 %)		2 (20 %)	
	worsened	1 (3.3 %)	1 (5 %)		–	

OC optic canal; ON optic nerve; ICA internal carotid artery