Erratum to: Endoscopic endonasal trans-sphenoidal approach for pituitary adenomas: Is one nostril enough?

Sheng Han · Xiaoxu Ding · Xinxin Tie · Yuan Liu ·
Junze Xia · Aihui Yan · Anhua Wu

Published online: 24 September 2013
© Springer-Verlag Wien 2013

Erratum to: Acta Neurochir (2013) 155:1601–1609
DOI: 10.1007/s00701-013-1788-8

The original version of this article unfortunately contained mistakes. On page 1607 and 1608, the figure legends for Fig 5 and 6 should be interchanged. Figure 5 shows images obtained from a 64-year-old female with a nonfunctioning macroadenoma who underwent the two-nostril approach. Figure 6 comprises images obtained from a 51-year-old man with a GH-secreting macroadenoma who underwent the one-nostril approach.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00701-013-1788-8>.

S. Han · X. Tie · Y. Liu · J. Xia · A. Wu (✉)
Department of Neurosurgery, The First Hospital of China,
Medical University, Nanjing Street 155, Heping District,
Shenyang 110001, China
e-mail: cmuwuanhua@yahoo.com.cn

X. Ding · A. Yan
Department of Otorhinolaryngology, The First Hospital of China,
Medical University, Shenyang, China