ERRATUM

Erratum to: Head injuries and the risk of concurrent cervical spine fractures

Tuomo Thesleff¹ · Anneli Kataja² · Juha Öhman¹ · Teemu M. Luoto¹

Published online: 23 March 2017
© Springer-Verlag Wien 2017

Erratum to: Acta Neurochir
DOI 10.1007/s00701-017-3133-0

Reason: Minor errors in Table 3 and related text “Patients with positive head CT scans (2.2%, n=6, did not have more neurosurgical operations within 1 year post-HI due to cervical spine fractures compared with patients with negative head CT scans (2.8%, n=23; p=0.614).”

Corrected version: “Patients with positive head CT scans (2.2%, n=6) did not have more neurosurgical operations within one year post-HI due to CSIs compared with patients with negative head CT scans (3.2%, n=26; p=0.431).”

The correct Table 3 is presented below:

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00701-017-3133-0>

✉ Tuomo Thesleff
tuomo.thesleff@pshp.fi

Anneli Kataja
anneli.kataja@pshp.fi

Juha Öhman
juha.ohman@pshp.fi

Teemu M. Luoto
teemu.luoto@pshp.fi

¹ Department of Neuroscience and Rehabilitation, Tampere University Hospital, P.O. Box 2000, FI-33521 Tampere, Finland

² Medical Imaging Centre, Department of Radiology, Tampere, University Hospital, P.O. Box 2000, FI-33521 Tampere, Finland

Table 3 The CSI - characteristics of the study sample stratified into subgroups (head CT-positive patients, head CT-negative patients, and whole sample)

Variable	Head CT-positive patients (<i>n</i> = 269)	%	Head CT-negative patients (<i>n</i> = 822)	%	<i>p</i> -value	Whole sample (<i>n</i> = 1091)	%
Cervical spine surgery within one year post injury	6	2.2	26	3.2	0.431	32	2.9
Cervical spine fracture	25	9.3	47	5.7	0.04	72	6.6
Cervical dislocation / subluxation	8	3.0	23	2.8	0.880	31	2.8
Number of fractured cervical vertebrae	35	13.0	66	8.0	0.014	101	9.3
C0	2	0.7	2	0.2	0.431	4	0.4
C1	5	1.9	5	0.6	0.062	10	0.9
C2 (odontoid)	5	1.9	8	1.0	0.245	13	1.2
C2 (non-odontoid)	5	1.9	11	1.3	0.538	16	1.5
C3	2	0.7	4	0.5	0.621	6	0.5
C4	1	0.4	8	1.0	0.344	9	0.8
C5	4	1.5	8	1.0	0.483	12	1.1
C6	8	3.0	9	1.1	0.031	17	1.6
C7	5	1.9	15	1.8	0.971	20	1.8
Multilevel cervical spine fracture (> 1 levels)	9	3.3	13	1.6	0.074	22	2.0
Patients with lower level cervical fracture (C3-C7)	16	5.9	31	3.8	0.127	47	4.3
Patients with upper level cervical fracture (C0-C2)	11	4.1	19	2.3	0.122	30	2.7
Spinal cord injury	1	0.4	12	1.5		13	1.2
ASIA-scale					0.490		
A	1	0.4	2	0.2		3	0.3
B	0	0	4	0.5		4	0.4
C	0	0	5	0.6		5	0.5
D	0	0	1	0.1		1	0.1
E or unknown	268	99.6	810	98.5		1078	98.8