Revisiting the taxonomy of the family *Circoviridae*: establishment of the genus *Cyclovirus* and removal of the genus *Gyrovirus*

Karyna Rosario¹ · Mya Breitbart¹ · Balázs Harrach² · Joaquim Segalés^{3,4} · Eric Delwart^{5,6} · Philippe Biagini⁷ · Arvind Varsani^{8,9}

Received: 9 December 2016 / Accepted: 29 December 2016 / Published online: 2 February 2017
© Springer-Verlag Wien 2017

Abstract The family *Circoviridae* contains viruses with covalently closed, circular, single-stranded DNA (ssDNA) genomes, including the smallest known autonomously replicating, capsid-encoding animal pathogens. Members of this family are known to cause fatal diseases in birds and pigs and have been historically classified in one of two genera: *Circovirus*, which contains avian and porcine pathogens, and *Gyrovirus*, which includes a single species (*Chicken anemia virus*). However, over the course of the past six years, viral metagenomic approaches as well as degenerate PCR detection in unconventional hosts and environmental samples have elucidated a broader host range, including fish, a diversity of mammals, and invertebrates, for members of the family *Circoviridae*. Notably, these methods have uncovered a distinct group of viruses that are closely related to members of the genus

Circovirus and comprise a new genus, *Cyclovirus*. The discovery of new viruses and a re-evaluation of genomic features that characterize members of the *Circoviridae* prompted a revision of the classification criteria used for this family of animal viruses. Here we provide details on an updated *Circoviridae* taxonomy ratified by the International Committee on the Taxonomy of Viruses in 2016, which establishes the genus *Cyclovirus* and reassigns the genus *Gyrovirus* to the family *Anelloviridae*, a separate lineage of animal viruses that also contains circular ssDNA genomes. In addition, we provide a new species demarcation threshold of 80% genome-wide pairwise identity for members of the family *Circoviridae*, based on pairwise identity distribution analysis, and list guidelines to distinguish between members of this family and other eukaryotic viruses with circular, ssDNA genomes.

Electronic supplementary material The online version of this article (doi:[10.1007/s00705-017-3247-y](https://doi.org/10.1007/s00705-017-3247-y)) contains supplementary material, which is available to authorized users.

✉ Karyna Rosario
krosari2@mail.usf.edu

✉ Arvind Varsani
arvind.varsani@asu.edu

¹ College of Marine Science, University of South Florida, Saint Petersburg, FL 33701, USA

² Institute for Veterinary Medical Research, Centre for Agricultural Research, Hungarian Academy of Sciences, Budapest, Hungary

³ Departament de Sanitat i Anatomia Animals, Universitat Autònoma de Barcelona, Bellaterra, 08193 Barcelona, Spain

⁴ UAB, Centre de Recerca en Sanitat Animal (CRESA, IRTA-UAB), Campus de la Universitat Autònoma de Barcelona, Bellaterra, 08193 Barcelona, Spain

⁵ Blood Systems Research Institute, San Francisco, California, USA

⁶ Department of Laboratory Medicine, University of California, San Francisco, San Francisco, California, USA

⁷ Viral Emergence and Co-evolution Unit, ADES, UMR 7268, Aix-Marseille University, CNRS, EFS, 27 Bd. Jean Moulin, 13005 Marseille, France

⁸ The Biodesign Center for Fundamental and Applied Microbiomics, Center for Evolution and Medicine, School of Life sciences, Arizona State University, Tempe, AZ 85287-5001, USA

⁹ Structural Biology Research Unit, Department of Clinical Laboratory Sciences, University of Cape Town, Observatory, Cape Town, South Africa

Introduction

The family *Circoviridae* was established in the mid-1990s when it was recognized that animal viruses with circular, single-stranded DNA (ssDNA) genomes were distinct from other eukaryotic ssDNA viruses classified at the time, including plant viruses with circular genomes (*Geminiviridae*) and animal viruses with linear genomes (*Parvoviridae*) [47, 83]. Originally, all the known animal viruses with covalently closed circular ssDNA genomes were classified in a single genus, *Circovirus*, within this family [47]. These animal viruses included avian and swine pathogens, namely beak and feather disease virus (BFDV), chicken anemia virus (CAV) and porcine circovirus (PCV, specifically the virus currently known as PCV-1). However, it is now recognized that animal viruses with circular ssDNA genomes are highly diverse and genome structure alone cannot be used for their taxonomical classification.

The last published report from the International Committee on Taxonomy of Viruses (ICTV), the 9th report reflecting the taxonomy from 2009, lists 12 viral species in the family *Circoviridae* [4]. These species are divided into two genera: *Circovirus*, which contained 11 species infecting birds and pigs, and *Gyrovirus*, which only included CAV. However, the progressive identification of new genomic sequences similar to members of the genus *Circovirus* in various animals and environmental samples subsequent to the release of the 9th report has prompted a revision of the classification framework used for this family. Furthermore, structural and genomic data indicate that CAV may represent a different lineage of ssDNA viruses and, thus, the genus *Gyrovirus* needed to be removed from the family *Circoviridae*. The genomic features of CAV are more closely aligned with ssDNA viruses in the family *Anelloviridae* and, thus, the genus *Gyrovirus* was reassigned to this family.

In this article we re-visit genomic features that characterize members of the family *Circoviridae* and provide an update to the taxonomy reported in the ICTV 9th report by: 1) establishing a new genus, *Cyclovirus*, that accommodates a distinct group of viruses closely related to members of the genus *Circovirus*; 2) reassigning the genus *Gyrovirus* to the family *Anelloviridae*; and 3) redefining the species demarcation criteria for members of the family *Circoviridae* and implementing a genome-wide pairwise identity based approach to classify known as well as new viruses that have been reported within the past six years.

Conserved genomic features among members of the family *Circoviridae*

Genus *Circovirus*

Most of what is known about circoviruses, or members of the genus *Circovirus*, comes from veterinary science-related research since these viruses are responsible for fatal diseases that affect birds (e.g., BFDV) and swine (e.g., PCV-2) [83]. In fact, until 2010, pigs were the only mammals known to be affected by circoviruses and most of the diversity for this group of viruses was reported from avian species (Table 1). However, studies employing viral metagenomic-based strategies and degenerate PCR for circoviruses in unconventional hosts have since identified the presence of circovirus genomes in freshwater fish [45, 46] and various mammals, including bats [34–36, 42, 91], chimpanzees [34], dogs [37], humans [34] and minks [41]. Although a definitive host has not been confirmed for some of these newly-detected circoviruses (e.g., bat-associated circoviruses), phylogenetic analyses indicate that circovirus genomes detected in mammals, in general, are more closely-related to each other than to avian circoviruses (Fig. 1).

Circovirus genomes range in size from ~1.8 to ~2.1 kb and are packaged within non-enveloped virions that have an icosahedral T = 1 structure and have an average diameter of ~15–25 nm [10, 69, 82, 83]. All circovirus genomes have an ambisense organisation containing open reading frames (ORFs) arranged on different strands of a dsDNA replicative form. Two major ORFs (>600 nt), encoding the replication-associated protein (Rep) on the virion strand and capsid protein (Cp) on the complementary strand of the replicative form, can be readily identified in circovirus genomes [4] (Fig. 2). The Rep, which has sequence motifs characteristic of proteins involved in rolling circle replication (RCR; see below) [28], is the most conserved circovirus protein. On the other hand, the Cp is significantly divergent and is only characterized by an N-terminal region rich in basic amino acids that may provide DNA binding activity [57]. Although Rep- and Cp-encoding ORFs are present in all circovirus genomes, other proteins may also be expressed by several circovirus species. For example, more than six ORFs have been identified in both avian and porcine circovirus genomes (e.g., [2, 23]) and BFDV virions have been consistently found associated with up to three proteins [68, 69]. Notably, porcine circoviruses (PCV-1 and PCV-2) are known to encode a third protein, denominated VP3, with apoptotic capacity [26, 32, 44] as well as a fourth one, ORF4, with a potential anti-apoptotic function [48].

Table 1 List of viral species classified within the family *Circoviridae*

Genus	Species/virus name	Acronym	Accession #	Isolation source (organism)	Isolation source (tissue)	Country	Reference
<i>Circovirus</i>	Barbel circovirus	BarCV	GU799606	<i>Barbus barbus</i>	Whole specimens (Fry stage)	Hungary	[45]
	Bat associated circovirus 1	BatACV-1	JX863737	<i>Rhinolophus ferrumequinum</i>	Guts	Myanmar	[24]
	Bat associated circovirus 2	BatACV-2	KC339249	<i>Rhinolophus ferrumequinum</i>	Guts	Myanmar	[24]
	Bat associated circovirus 3	BatACV-3	JQ814849	<i>Rhinolophus ferrumequinum</i>	Pharyngeal & rectal swabs	China	[90]
	Bat associated circovirus 4	BatACV-4	KT783484	<i>Tadarida brasiliensis</i>	Pharyngeal swabs	Brazil	[42]
	Bat associated circovirus 5	BatACV-5	KJ641727	<i>Plecotus auritus</i>	Pharyngeal & rectal swabs	China	[91]
	Bat associated circovirus 6	BataCV-6	KJ641724	<i>Rhinolophus affinis</i>	Pharyngeal & rectal swabs	China	[91]
	Bat associated circovirus 7	BatACV-7	KJ641723	<i>Rhinolophus sinicus</i>	Pharyngeal & rectal swabs	China	[91]
	Bat associated circovirus 8	BatACV-8	KJ641711	<i>Myotis ricketti</i>	Pharyngeal & rectal swabs	China	[91]
	Beak and feather disease virus*	BFDV	AF071878	Psittaciformes species	Skin	USA	[57]
	Canary circovirus*	CaCV	AJ301633	<i>Serinus canaria</i>	Organs	Italy	[84]
	Canine circovirus	CanineCV	KC241982	<i>Canis lupus familiaris</i>	Liver	USA	[37]
	Chimpanzee associated circovirus 1	ChimpACV-1	GQ404851	<i>Pan troglodytes</i>	Faeces	Rwanda	[34]
	Duck circovirus*	DuCV	DQ100076	<i>Anas domesticus</i>	Liver, spleen, thymus, & <i>Bursa fabricii</i>	USA	[1]
	European catfish circovirus	EcatfishCV	JQ011377	<i>Silurus glanis</i>	Liver, spleen, gills, kidneys & gonads	Hungary	[46]
	Finch circovirus*	FiCV	DQ845075	<i>Chloebia gouldiae</i>	<i>Bursa fabricii</i>	N/A	[86]
	Goose circovirus*	GoCV	AJ304456	Goose	<i>Bursa fabricii</i>	Germany	[85]
	Gull circovirus*	GuCV	DQ845074	<i>Larus argentatus</i>	<i>Bursa fabricii</i>	Sweden	[86]
	Human associated circovirus 1	HuACV-1	GQ404856	<i>Homo sapiens</i>	Faeces	Nigeria	[34]
	Mink circovirus	MiCV	KJ020099	<i>Mustela sp.</i>	Gut	China	[41]
	Pigeon circovirus*	PiCV	AF252610	<i>Columbia livia</i>	<i>Bursa fabricii</i>	Germany	[52]
	Porcine circovirus 1*	PCV-1	AF071879	N/A	Cell culture	N/A	[57]
	Porcine circovirus 2*	PCV-2	AF027217	<i>Sus domesticus</i>	Lungs, lymph nodes, spleens, & tonsils	Canada	[23]
	Raven circovirus*	RaCV	DQ146997	<i>Corvus coronoides</i>	Feathers	Australia	[78]
	Starling circovirus*	StCV	DQ172906	<i>Sturnus vulgaris</i>	Spleen	Spain	[29]
	Swan circovirus*	SwCV	EU056309	<i>Cygnus olor</i>	Liver and spleen	Germany	[22]
	Zebra finch circovirus	ZfiCV	KP793918	<i>Taeniopygia guttata</i>	N/A	Germany	[67]
<i>Cyclovirus</i>	Bat associated cyclovirus 1	BatACyV-1	HM228874	<i>Antrozous pallidus</i>	Faeces	USA	[35]
	Bat associated cyclovirus 2	BatACyV-2	JF938079	<i>Myotis spp.</i>	Faeces	China	[18]
	Bat associated cyclovirus 3	BatACyV-3	JF938081	<i>Myotis spp.</i>	Faeces	China	[18]
	Bat associated cyclovirus 4	BatACyV-4	JF938082	<i>Myotis spp.</i>	Faeces	China	[18]

Table 1 continued

Genus	Species/virus name	Acronym	Accession #	Isolation source (organism)	Isolation source (tissue)	Country	Reference
	Bat associated cyclovirus 5	BatACyV-5	HQ738637	<i>Tadarida brasiliensis</i>	Muscle	USA	[36]
	Bat associated cyclovirus 6	BatACyV-6	KJ641712	<i>Rhinolophus pusillus</i>	Pharyngeal & rectal swabs	China	[91]
	Bat associated cyclovirus 7	BatACyV-7	KJ641740	<i>Rhinolophus pusillus</i>	Pharyngeal & rectal swabs	China	[91]
	Bat associated cyclovirus 8	BatACyV-8	KJ641715	<i>Rhinolophus pusillus</i>	Pharyngeal & rectal swabs	China	[91]
	Bat associated cyclovirus 9	BatACyV-9	KJ641720	<i>Tylonycteris pachypus</i>	Pharyngeal & rectal swabs	China	[91]
	Bat associated cyclovirus 10	BatACyV-10	KM382270	<i>Molossus molossus</i> , <i>Tadarida brasiliensis</i>	Faeces	Brazil	[43]
	Bat associated cyclovirus 11	BatACyV-11	KJ641717	<i>Myotis spp.</i>	Pharyngeal & rectal swabs	China	[91]
	Bat associated cyclovirus 12	BatACyV-12	KM382269	<i>Molossus molossus</i> , <i>Tadarida brasiliensis</i>	Faeces	Brazil	[43]
	Bat associated cyclovirus 13	BatACyV-13	KJ641728	<i>Plecotus auritus</i>	Pharyngeal & rectal swabs	China	[91]
	Bat associated cyclovirus 14	BatACyV-14	KT732785	<i>Pteropus tonganus</i>	Faeces	Tonga	[49]
	Bat associated cyclovirus 15	BatACyV-15	KT732786	<i>Pteropus tonganus</i>	Faeces	Tonga	[49]
	Bat associated cyclovirus 16	BatACyV-16	KT732787	<i>Pteropus tonganus</i>	Faeces	Tonga	[49]
	Bovine associated cyclovirus 1	BoACyV-1	HQ738634	<i>Bos taurus</i>	Muscle	Pakistan	[36]
	Chicken associated cyclovirus 1	ChickACyV-1	HQ738643	<i>Gallus gallus</i>	Muscle	Nigeria	[36]
	Chimpanzee associated cyclovirus 1	ChimpACyV-1	GQ404849	<i>Pan troglodytes</i>	Faeces	Central Africa	[34]
	Cockroach associated cyclovirus 1	CroACyV-1	JX569794	<i>Eurycotis floridana</i>	Abdomen	USA	[59]
	Dragonfly associated cyclovirus 1	DfACyV-1	JX185419	<i>Pantala flavescens</i>	Abdomen	Tonga	[70]
	Dragonfly associated cyclovirus 2	DfACyV-2	JX185422	<i>Pantala flavescens</i>	Abdomen	USA	[71]
	Dragonfly associated cyclovirus 3	DfACyV-3	JX185424	<i>Erythemis simplicicollis</i>	Abdomen	USA	[71]
	Dragonfly associated cyclovirus 4	DfACyV-4	KC512916	<i>Aeshna multicolor</i>	Abdomen	USA	[11]
	Dragonfly associated cyclovirus 5	DfACyV-5	JX185426	<i>Erythrodiplax umbrata</i>	Abdomen	Puerto Rico	[71]
	Dragonfly associated cyclovirus 6	DfACyV-6	KC512918	<i>Aeshna multicolor</i>	Abdomen	USA	[11]
	Dragonfly associated cyclovirus 7	DfACyV-7	KC512919	<i>Xanthocnemis zealandica</i>	Abdomen	New Zealand	[11]
	Dragonfly associated cyclovirus 8	DfACyV-8	KC512920	<i>Orthetrum Sabina</i>	Abdomen	Australia	[11]
	Feline associated cyclovirus 1	FeACyV-1	KM017740	<i>Felis catus</i>	Faeces	USA	[92]
	Goat associated cyclovirus 1	GoACyV-1	HQ738636	<i>Capra aegagrus hircus</i>	Muscle	Pakistan	[36]

Table 1 continued

Genus	Species/virus name	Acronym	Accession #	Isolation source (organism)	Isolation source (tissue)	Country	Reference
	Horse associated cyclovirus 1	HoACyV-1	KR902499	<i>Equus caballus</i>	Nasal secretions	USA	[38]
	Human associated cyclovirus 1	HuACyV-1	GQ404847	<i>Homo sapiens</i>	Faeces	Pakistan	[34]
	Human associated cyclovirus 2	HuACyV-2	GQ404844	<i>Homo sapiens</i>	Faeces	Pakistan	[34]
	Human associated cyclovirus 3	HuACyV-3	GQ404846	<i>Homo sapiens</i>	Faeces	Pakistan	[34]
	Human associated cyclovirus 4	HuACyV-4	GQ404857	<i>Homo sapiens</i>	Faeces	Tunisia	[34]
	Human associated cyclovirus 5	HuACyV-5	GQ404845	<i>Homo sapiens</i>	Faeces	Pakistan	[34]
	Human associated cyclovirus 6	HuACyV-6	GQ404854	<i>Homo sapiens</i>	Faeces	Nigeria	[34]
	Human associated cyclovirus 7	HuACyV-7	GQ404855	<i>Homo sapiens</i>	Faeces	Nigeria	[34]
	Human associated cyclovirus 8	HuACyV-8	KF031466	<i>Homo sapiens</i>	Cerebrospinal fluid	Vietnam	[80]
	Human associated cyclovirus 9	HuACyV-9	KC771281	<i>Homo sapiens</i>	Blood serum	Malawi	[76]
	Human associated cyclovirus 10	HuACyV-10	KF726984	<i>Homo sapiens</i>	Respiratory secretion	Chile	[63]
	Human associated cyclovirus 11	HuACyV-11	KJ831064	<i>Homo sapiens</i>	Cerebrospinal fluid	Sri Lanka	[64]
	Squirrel associated cyclovirus 1	SqACyV-1	LC018134	<i>Callosciurus erythraeus taiwanensis</i>	Stomach contents	Japan	[73]

Viral species described before 2010 are marked with an asterisk

N/A = information not available

The presence of a well-conserved Rep in circovirus genomes suggests that these viruses replicate through RCR and recombinant expression studies support this idea [7, 16, 51, 77]. Circovirus Reps are characterized by the presence of three RCR motifs at the N-terminus including, motif I [FT(L/I)NN], motif II [PHLQG] and motif III [YC(S/x)K] where “x” represents any residue [72] (Fig. 3). In addition to the RCR domain, circovirus Reps contain dNTP-binding or P-loop NTPase domains characteristic of superfamily 3 (SF3) helicases which are distinguished by the presence of three conserved motifs tightly packed in a domain containing ~100 amino acids [19, 20]. The SF3 helicase motifs found in circovirus Reps include Walker-A [G(P/x)(P/x)GxGK(S/t)], Walker-B [uuDDF], and motif C [uTSN] where “x” represents any residue, “u” represents a hydrophobic amino acid (i.e., F, I, L, V, M), and residues in lower case are observed at lower frequency [72].

In addition to conserved gene synteny and the presence of a Rep with characteristic RCR and helicase motifs, circovirus genomes have two intergenic regions (IR) and a

conserved origin of replication (*ori*) (Fig. 2). The *ori* is located within the IR located between the 5' ends of the Rep- and Cp-encoding ORFs [50]. The circovirus *ori* is characterized by a conserved nonanucleotide motif ‘(T/n)A(G/t)TATTAC’ (Fig. 2), where lower case nucleotides are observed at low frequency and ‘n’ represents any nucleotide, located at the apex of a potential stem-loop structure [50, 72]. The Rep introduces a nick in the virion-sense strand between positions 7 and 8 of the nonanucleotide motif (i.e., TAGTATT¹AC), presumably initiating circovirus genome replication through RCR [77].

Genus *Cyclovirus*, a newly established taxon

In 2010, a group of viruses most-closely related to circoviruses were discovered through viral metagenomic analysis and degenerate PCR in stool samples from primates (humans and chimpanzees) and meat products from various animals (camels, chickens, cows, goats, and sheep) [34]. These viruses were tentatively named cycloviruses to distinguish them from the already known circoviruses,

Circovirus *Cyclovirus*

Fig. 1 Maximum likelihood (ML) phylogenetic trees of representative sequences from members of viral species within the genus *Circovirus* (top) and the genus *Cyclovirus* (bottom). The ML trees were inferred using PhyML [21] with the GTR+G model of substitution after aligning complete genome sequences using the MUSCLE algorithm [15]. Branches with <80% SH-like support have been collapsed. The phylogenetic trees of circoviruses and cycloviruses were rooted after using cyclovirus or circovirus reverse complemented genome sequences as an outgroup, respectively. Species acronyms are defined in Table 1

while still highlighting the circular topology of their genomes [34]. Cycloviruses are closely related to circoviruses and share genomic features with this well-known group of animal pathogens [14]. However, phylogenetic and genomic differences between circoviruses and cycloviruses

justified the creation of a second taxonomic unit, the newly established genus *Cyclovirus*, within the family *Circoviridae*.

In contrast to circoviruses, cycloviruses have been found associated with both vertebrates and invertebrates. Although cycloviruses were discovered in stool samples from primates [34], cyclovirus genomes have now been reported from a diversity of specimens including mammals (bats, cats, cows, goats, horses, squirrels, sheep) [18, 34, 36, 38, 43, 49, 73, 91, 92], birds (chickens) [34, 36], and insects (cockroaches and dragonflies) [11, 59, 70, 71] (Table 1). Additionally, a diversity of cyclovirus genomes have been recovered from human samples other than faeces [17, 34, 80], including cerebrospinal fluid [80], blood serum [76], and respiratory

Fig. 2 Genome schematics illustrate the major open reading frames (ORFs) characteristic of members of the *Circoviridae* (left) and *Anelloviridae* families (right). Members of the family *Circoviridae*, including the *Circovirus* and *Cyclovirus* genera, have two major ORFs encoding replication-associated (Rep) and capsid (Cp) proteins as well as a conserved nonanucleotide motif marking the origin of replication. The nonanucleotide motif sequence is depicted through sequence probability logos generated in Weblogo 3 [9]. Note that the orientation of major ORFs relative to the nonanucleotide motif differs between genomes representing the *Circovirus* and *Cyclovirus* genera. The *rep* of the *Cyclovirus* type species, Human associated cyclovirus 8 (HuACyV-8), is interrupted by an intron. Although the presence of

introns has been observed in various cyclovirus genomes, this has not been reported for circoviruses (Supplemental Figures 1 and 2). In contrast to the *Circoviridae*, members of the family *Anelloviridae* consistently have three major ORFs in an unisense organization. The genera *Gyrovirus*, which currently includes Chicken anemia virus (CAV) alone, and *Alphatorquevirus*, with Torque teno virus 1 (TTV-1) as the type species, are shown to highlight similarities among these ssDNA viruses. The CAV-VP1 and TTV-1 ORF1 are thought to represent capsid proteins. Proteins labelled as VP2 and ORF2 in CAV and TTV-1 genome schematics, respectively, contain a motif characteristic of protein tyrosine phosphatases. In addition, the CAV-VP3 and TTV-1 ORF3 proteins exhibit apoptotic activity

Fig. 3 Sequence probability logos generated using Weblogo 3 [9] highlighting conserved amino acid residues characteristic of rolling circle replication (RCR), including RCR motifs I through III, and superfamily 3 (SF3) helicase motifs, including Walker A and B as well as motif C, found in replication-associated proteins (Rep). The logos were generated using representative sequences from members

of the 70 species in the family *Circoviridae* (*Circovirus*, n=27; *Cyclovirus*, n=43). Numbers below *Circovirus* and *Cyclovirus* logos indicate the relative amino acid position for each motif based on the type species, including Porcine circovirus 1 (accession number AF071879) and Human associated cyclovirus 8 (accession number KF031466), respectively

secretions [63]. However, it has been difficult to identify a definitive host for most, if not all, cycloviruses since these viruses have only been identified through metagenomic analysis and degenerate PCR. This is further complicated by the fact that many cycloviruses have been discovered from guts and fecal samples (Table 1), which may include viruses that are dietary in origin. Moreover, phylogenetic analysis of cyclovirus genomes did not reveal any clusters by the type of organism from which they were identified (Fig. 1).

Similar to circoviruses, cycloviruses have small genomes (~1.7 to 1.9 kb) that contain two major ORFs encoding the Rep and Cp organized in two different strands of a dsDNA form (Fig. 2) [34, 72]. Both Rep and Cp of cycloviruses have similar features when compared with their corresponding circovirus proteins (reviewed by [14, 72]). The cyclovirus Rep contains motifs similar to circovirus RCR and SF3 helicase motifs, including RCR motifs I [FT(L/W)NN], II [(P/x)HLQG] and III [Y(C/I)(S/x)K] and SF3 helicase motifs Walker-A [G(P/x)(P/t)(G/x)xGKS], Walker-B [uuDDF], and motif C [uTS(N/e)] where “x” represents any type of residue and “u” represents a hydrophobic amino acid (i.e., F, I, L, V, M). The

cyclovirus putative Cp is also characterized by an N-terminal region rich in basic amino acids as seen in circovirus Cps. Furthermore, the putative cyclovirus *ori* is marked by the same conserved nonamer observed in circoviruses (‘TAGTATTAC’) located at the apex of a potential stem-loop structure (Fig. 2) [72].

Despite these similarities, phylogenetic analyses of the Rep and Cp show that circoviruses and cycloviruses form distinct clades (Fig. 4). Furthermore, there are key genomic features that distinguish cycloviruses from circoviruses (Fig. 2). Cyclovirus genomes contain an IR between the 5' ends of Rep- and Cp-encoding ORFs; however, the IR between the 3' ends of these major ORFs is either absent or consistently smaller than that observed in circovirus genomes [34, 72]. Additionally, cycloviruses and circoviruses might employ slightly different replication and transcription strategies since their genome coding regions are organized differently. Although the cyclovirus virion strand has not been empirically determined, it is predicted that these viruses encapsidate the strand containing the conserved nonanucleotide motif based on what has been shown for other eukaryotic Rep-encoding, ssDNA viruses [72]. Using the nonanucleotide motif as a point of

Fig. 4 Maximum likelihood (ML) phylogenetic trees of representative replication-associated protein (Rep; left) rooted with Reps of closely related sequences (GenBank accession #s JX904473, KC248418 and KJ547623) and mid-point rooted capsid protein (Cp; right) amino acid sequences of representatives from viral species

reference, it is clear that the genome organization of cycloviruses seems to be a mirror image from the one seen in circoviruses. The putative *ori*, which is marked by the nonanucleotide motif, is located on the Rep-encoding strand of circoviruses, while in cycloviruses is located on the Cp-encoding strand [72]. In addition to differences in genomic arrangement, the presence of introns has been identified within the ORFs of several cyclovirus genomes, whereas this has not been reported for circoviruses (Supplemental Figures 1 and 2).

Reassignment of the genus *Gyrovirus* to the family *Anelloviridae*

Since CAV, BFDV, and PCV were the first described animal viruses with circular ssDNA genomes, studies sought to better understand the relationship between these

within the *Circovirus* (blue font) and *Cyclovirus* (purple font) genera. The ML trees were inferred using PhyML [21] with the LG model of substitution after aligning amino acid sequences using the MUSCLE algorithm [15]. Branches with <80% SH-like support have been collapsed. Species acronyms are defined in Table 1

three viruses. As early as 1991 it was noted that CAV virions had marked physicochemical and morphological differences when compared to BFDV and PCV, which were indistinguishable under the same conditions [82]. Therefore, even before animal viruses with circular ssDNA genomes were classified, it was 'inadvisable' to place CAV in the same family as BFDV and PCV based on comparative electron microscopy [82]. Despite these observations, CAV became the type species of the family *Circoviridae* when it was originally reported as an official virus family in 1995 with the release of the ICTV 6th Report [47]. Nevertheless, early on it was recognized that CAV did not share the same characteristics as BFDV and PCV and thus a second genus, *Gyrovirus*, was created within the family *Circoviridae* in 1999 to accommodate CAV [27]. The genus *Gyrovirus*, with CAV as its sole member, remained in the family for more than a decade. However, mounting genomic, molecular, and structural data indicated that CAV

represents a different lineage of ssDNA viruses as originally suspected, thus granting the placement of the genus *Gyrovirus* in a separate family.

Although CAV virions are non-enveloped and icosahedral, they are larger than circovirus virions and have a unique structure with protruding pentagonal shaped units compared to flat pentameric units observed in circoviruses [10, 82]. In addition to appearing structurally unrelated to circoviruses, CAV has a genomic architecture that is radically different from that of circoviruses. The CAV genome has a negative-sense organization, containing three major overlapping ORFs in the virion or genomic strand [31, 58] (Fig. 2). Moreover, CAV genomes do not encode an identifiable Rep and lack the conserved nonanucleotide motif that marks the *ori* of circoviruses and cycloviruses [53, 57, 72].

Gyroviruses do not appear to be either structurally or genetically related to members of the family *Circoviridae*. Instead, the genomic features of CAV are reminiscent of members of the family *Anelloviridae* (Fig. 2). The family *Anelloviridae*, which accommodates 12 genera of vertebrate-infecting viruses, currently represents the only other family of animal viruses with covalently closed, circular ssDNA genomes [5]. Although sequence variability within anelloviruses is extremely high, both at the nucleotide and amino acid levels, their genomes have common features [3]. It is important to note that the common genomic features that justified the creation of the family *Anelloviridae* [3] are also observed in CAV genomes. Similar to CAV, anellovirus genomes have a negative-sense organization with overlapping ORFs with similar relative sizes [3, 12, 54, 79]. In addition, anellovirus and CAV genomes contain GC-rich stretches in non-coding regions where replication is thought to initiate due to the presence of potential stem-loop structures [2, 13]. The non-coding or untranslated region of CAV and some anellovirus genomes also contains similar regulatory elements and share high identity within a 36-nt stretch [25, 54]. Moreover, CAV and some anelloviruses have similar transcription patterns [25, 30].

Although CAV and anelloviruses do not have significant sequence similarities, these viruses may encode proteins with similar functions. The largest ORF observed in CAV and anellovirus genomes encodes for a putative structural protein, named VP1 and ORF1, respectively. Both VP1 and ORF1 are characterized by an N-terminal region rich in basic amino acids, which resembles Cps encoded by members of the family *Circoviridae* [3, 57]. Biochemical characterization of CAV virions revealed VP1 as the sole Cp component for this gyrovirus [81]; however, this has not been conclusively demonstrated for the anellovirus ORF1. CAV and anellovirus genomes also have two other ORFs that encode proteins with similar characteristics and

are of comparable sizes. The VP2 and ORF2 of CAV and anellovirus genomes, respectively, encode a protein containing a motif characteristic of protein tyrosine phosphatases (i.e., $WX_7HX_3CXCX_5H$, where 'X' represents any residue) [60]. In addition, the CAV-VP3 and ORF3 of some anelloviruses encode a protein with comparable apoptotic capacity [33, 65].

Based on similarities in genome architectures and transcription profiles, gyroviruses are more closely related to members of the family *Anelloviridae* rather than to those of the *Circoviridae*. Therefore, the genus *Gyrovirus* has been reassigned to the former family. Until 2011, CAV was the only described gyrovirus species. However, other potential members of the genus *Gyrovirus* have been recently discovered in humans and birds. These novel gyroviruses (GyVs) have been identified in human skin (human gyrovirus (HGyV) [74]) and feces (GyV3 through GyV6; [8, 61, 62]), chicken serum (avian gyrovirus 2 (AGV2) [66]) and meat (GyV4 and GyV7; [8]) as well as spleen and uropygial gland tissues from sea birds (GyV8; [39]). Since HGyV and AGV2 may represent the same species [61], the genus *Gyrovirus* should soon be updated to reflect at least eight viral species.

Revised species demarcation criteria for the family *Circoviridae*

The ICTV 9th Report specified a species demarcation criteria for members of the family *Circoviridae* (i.e., genus *Circovirus*) of <75% genome-wide identity and <70% amino acid identity for the Cp [4]. However, these criteria were based on the distribution of pairwise identities derived from global alignments which may be inaccurate due to inconsistencies introduced during alignment and gap-handling issues [55]. Furthermore, the above mentioned species demarcation criteria did not include members of the newly established genus *Cyclovirus* and circovirus-related genomes that have been reported since 2009. Therefore, there is a need to re-evaluate the species demarcation criteria for members of the family *Circoviridae*. Here we use a pairwise identity distribution analysis to establish species demarcation criteria to classify viruses within the family *Circoviridae*.

Pairwise identities derived from aligning pairs of genome sequences individually have been recently used as a more standardized and accurate method to calculate genome identity scores [55, 56]. Hence, we reanalyzed and calculated genome-wide pairwise identities for all the circovirus and cyclovirus genomes available in public databases through June 2016 using SDT v1.2 [56] (Supplemental Figures 3 and 4; Supplemental File 1). The analysis shows that a pairwise identity species cut-off

between 78% - 80% is best suited for both circoviruses and cycloviruses (Fig. 5). Since PCV-1 and PCV-2 share 79% genome-wide pairwise identity (Supplemental Figure 3) and these viruses have been historically considered to represent separate species due to marked biological differences (i.e., pathogenicity), we have delineated the species demarcation threshold at 80%. Therefore, the newly established species demarcation criteria will maintain the current classification for all previously characterized circovirus species, including the closely-related PCV-1 and PCV-2. The 80% genome-wide pairwise identity cut-off is well supported phylogenetically for both circoviruses and cycloviruses (Fig. 1). As a general rule, viruses that have <80% genome-wide pairwise identities with other members of the family *Circoviridae* coupled with phylogenetic support should be considered distinct species.

The 80% identity threshold for distinct species generally holds true for pairwise comparisons of either the *cp* or *rep* gene sequences (Fig. 5). However, only complete genomes

should be considered for assignment of new species in the event that they share <80% pairwise identity with any classified species within the family *Circoviridae*. Notably, pairwise comparisons between circoviruses and cycloviruses indicate that members of the family *Circoviridae* can have overlapping pairwise identity ranges based on the Rep. Cyclovirus Reps may share up to 50% amino acid pairwise identity with those of some circoviruses, which is higher than pairwise identities observed among cycloviruses (as low as 36%) and circoviruses (as low as 40%) alone (Supplemental Figure 5). Therefore, the classification of newly identified viruses with significant similarities to either circoviruses or cycloviruses should be primarily based on genomic characteristics and genome-wide pairwise identities. Rep identities and genome organization (specifically the location of the *ori* relative to coding regions) should be used to identify a genome as belonging to either the *Circovirus* or *Cyclovirus* genus. Once this distinction is made, circovirus and cyclovirus nucleotide

Fig. 5 Distribution of pairwise identities among members of the genus *Cyclovirus* (purple bars; left) and the genus *Circovirus* (blue bars; right). Plots reflect pairwise identities based on calculations for complete genome sequences (top) as well as the replication-

associated (*rep*; middle) and capsid (*cp*; bottom) genes. All pairwise identities were calculated using the Sequence Demarcation Tool version 1.2 [56] with the MUSCLE alignment algorithm [15]

sequences should be analysed independently for pairwise identity comparisons and taxonomic classification. Otherwise, obtaining a reliable alignment of complete genome sequences is extremely difficult since cycloviruses and circoviruses have a different genome organization. Even if cyclovirus genomes could be aligned to circoviruses to some degree by reverse complementing cyclovirus sequences, such an approach is not recommended since the Cp is highly divergent compared to the Rep.

After determining that 80% genome-wide pairwise identity was a suitable threshold for species demarcation, we classified all of the *Circoviridae*-related genomes available in GenBank through the end of June 2016 (Table 1). Note that the original published names for many of the genomes have been changed for consistency and that viruses for which a definitive host has not been identified are denoted by the presence of the word ‘associated’ in the species name. The latest analysis indicates that there are 27 distinct circovirus species. PCV-2 is, by far, the species with the most reported genome sequences, followed by BFDV, reflecting the interest in these animal pathogens over the years (Fig. 6). After the ICTV ratification of circovirus species listed here, a third porcine circovirus species, named PCV-3, was discovered [40]. The PCV-3 genome sequence was not included in the analyses presented in this taxonomy update; however, preliminary analyses indicate that indeed PCV-3 represents a new circovirus species. Although cycloviruses were discovered more recently than circoviruses, there are twice as many species ($n = 43$) reported to date that belong to the *Cyclovirus* genus than there are species from the genus *Circovirus*. It is possible that cycloviruses are more diverse and widespread than circoviruses since the former have been reported from both vertebrate and invertebrate animals, whereas circovirus genomes have only been found associated with birds, freshwater fish, and mammals.

Recommendations for classifying *Circoviridae* sequences

As more metagenomic analyses are performed on a diversity of organisms and environments, it is expected that novel circoviruses and cycloviruses will continue to be identified. In an effort to standardize the taxonomy framework for the family *Circoviridae*, here we outline a few guidelines regarding how to analyse and report genomes representing members of this family.

- 1) **Genome sequence verification.** ICTV will now consider classifying viral species based on complete genome sequences derived from metagenomic analyses [75]. Since the genomes of members of the

Fig. 6 Diversity of variants within each assigned species in the *Circovirus* (blue font) and *Cyclovirus* (purple font) genera. The total number of isolates within the species is given in parentheses by each species name. Details regarding genome sequences used for this analysis are provided in Supplementary File 1

family *Circoviridae* are relatively small (~ 2 kb) and have a circular topology, efforts should be made to close genomes and, whenever possible, verify genomes through inverse PCR. One of the biggest limitations of metagenomically-derived genomes is that there are no isolates of the viruses being described. Amplifying complete genomes through inverse PCR and cloning before sequencing will provide a biological archive of described viral genomes within plasmids. Additionally, working with PCR-verified genomes will ensure that the reported sequences are not chimeric entities assembled from metagenomic data.

- 2) **Submitting sequences to GenBank.** It is important to provide the correct annotation for major proteins (Rep and Cp) when submitting genome sequences to GenBank, including the identification and removal of introns. In addition, for consistency, all genomes should be submitted starting with the first nucleotide of the nonanucleotide motif. This will ensure that the genomes are reported in the correct orientation, which is a key feature for distinguishing between circovirus and cyclovirus genomes. All species level representative sequences analysed for this manuscript have been provided as supplemental material following the format indicated above, including complete genomes (Supplemental Files 2A and 2B) as well as Rep (Supplemental Files 3A and 3B) and Cp (Supplemental Files 4A and 4B) coding sequences.
- 3) **Naming species.** The following basic rules should be implemented when naming viral species:
 - a. If a given genome has >80% genome-wide pairwise identity with a genome sequence from a member of a previously described viral species, the name from the existing species should be adopted and a specific isolate name should be provided. For example, a newly identified virus in foxes that has a genome-wide pairwise identity of >80% with sequences from members of the species *Canine circovirus* could be named ‘*Canine circovirus* [fox-associated isolate 1]’.
 - b. If a novel virus representing a new species is being described (i.e., the genome has <80% genome-wide pairwise identity with known viral sequences) the word ‘associated’ should be added as a modifier to the species name, unless

there is biological data identifying a definitive host (e.g., *Bat associated cyclovirus* # species).

- 4) **Resolving species assignment conflicts.** There might be instances when it is difficult to classify a new sequence based on the established species demarcation criteria. Here we provide guidelines on how to approach conflicts based on recommendations for classifying species from the family *Geminiviridae*, another group of Rep-encoding circular, ssDNA viruses [6, 55, 87, 88]. Species assignments might be uncertain when:

- a. A given sequence shares >80% genome-wide pairwise identity with viral sequences within a particular species, but <80% identity with other isolates of that same species. This will lead to more than 20% sequence divergence among variants of a single species (e.g., HuACyV-8, see Fig. 6). To resolve this conflict, the new sequence should be classified within any species with which it shares >80% identity based on a given classified isolate, even if it is <80% identical to other isolates within that species.
- b. A given sequence shares >80% genome-wide pairwise identity with viral sequences assigned to two or more different species. In such cases, it is suggested that the new sequence be considered as belonging to the species with which it shares the highest degree of similarity.

- 5) **Reporting non-*Circoviridae* genomes.** A growing number of circular Rep-encoding ssDNA (CRESS DNA) viruses have been reported from various organisms and environments [14, 72] and it is expected that more genomes will continue to be reported. Many of these novel CRESS DNA genome sequences have similarities with members of the family *Circoviridae* based on the Rep. It is important to note that only sequences that clearly represent members of the *Circovirus* or *Cyclovirus* genera, based on genome features discussed here, should be classified as part of the family *Circoviridae*. If novel sequences with best matches to circovirus or cyclovirus genomes in GenBank share <55% genome-wide pairwise identity (Supplemental Figures 3 - 4) with those sequences, they should be considered as unclassified CRESS DNA viral sequences for the time being. Although additional genera in the family *Circoviridae* may be created in the future, this level of diversity (i.e., 45% divergence) is similar to the diversity that has been observed in other CRESS DNA viral families, such as the *Geminiviridae* and *Genomoviridae* [89].

Concluding remarks

The growing number of genomes representing members of the family *Circoviridae* in the database required a re-evaluation of the taxonomy classification framework for this family. Here we have reviewed genomic features that characterize members of the family *Circoviridae*, outlined guidelines on how to analyse genome sequences, and provided a current list of circovirus and cyclovirus species based on the newly established species demarcation criteria. These efforts should facilitate future analyses geared towards elucidating evolutionary relationships among classified as well as newly identified members of the family *Circoviridae*.

Acknowledgements KR and MB are supported by the National Science Foundation's Assembling the Tree of Life Program Grant DEB-1239976.

Compliance with ethical standards

There are no conflicts of interest; the research did not involve human participants or animals. The data used for the analyses in this manuscript is publicly available in GenBank.

Disclaimer This article is based on the taxonomic proposals 2016.007a,bD.A.v2.Circovirus_5sp and 2016.008a,bD.A.v2.Cyclovirus_15sp which has been considered and approved by the Executive Committee (EC) of the ICTV. AV and BH are elected members of the ICTV EC. KR, MB, JS and ED are current members of the *Circoviridae* study group of ICTV, whereas PB is a past member of the *Circoviridae*-*Anelloviridae* group.

References

1. Banda A, Galloway-Haskins RI, Sandhu TS, Schat KA (2007) Genetic analysis of a duck circovirus detected in commercial Pekin ducks in New York. *Avian Dis* 51:90–95
2. Bassami MR, Berryman D, Wilcox GE, Raidal SR (1998) Psittacine beak and feather disease virus nucleotide sequence analysis and its relationship to porcine circovirus, plant circoviruses, and chicken anemia virus. *Virology* 249:453–459
3. Biagini P (2009) Classification of TTV and related viruses (anelloviruses). In: de Villiers E-M, Hausen Hz (eds) *TT viruses: the still elusive human pathogens*. Springer, Berlin, pp 21–33
4. Biagini P, Bendinelli M, Hino S, Kakkola L, Mankertz A, Niel C, Okamoto H, Raidal S, Teo CG, Todd D (2012) Family *Circoviridae*. In: King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ (eds) *Virus taxonomy: classification and nomenclature of viruses: ninth report of the International Committee on Taxonomy of Viruses*. Academic Press, London, pp 343–349
5. Biagini P, Bendinelli M, Hino S, Kakkola L, Mankertz A, Niel C, Okamoto H, Raidal S, Teo CG, Todd D (2012) Family *Anelloviridae*. In: King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ (eds) *Virus taxonomy: classification and nomenclature of viruses: ninth report of the International Committee on Taxonomy of Viruses*. Academic Press, London, pp 331–341

6. Brown J, Zerbini FM, Navas-Castillo J, Moriones E, Ramos-Sobrinho R, Silva JF, Fiallo-Olivé E, Briddon R, Hernández-Zepeda C, Idris A, Malathi VG, Martin D, Rivera-Bustamante R, Ueda S, Varsani A (2015) Revision of *Begomovirus* taxonomy based on pairwise sequence comparisons. *Arch Virol* 160:1593–1619
7. Cheung AK (2006) Rolling-circle replication of an animal circovirus genome in a theta-replicating bacterial plasmid in *Escherichia coli*. *J Virol* 80:8686–8694
8. Chu DKW, Poon LLM, Chiu SSS, Chan KH, Ng EM, Bauer I, Cheung TK, Ng IHY, Guan Y, Wang D, Peiris JSM (2012) Characterization of a novel gyrovirus in human stool and chicken meat. *J Clin Virol* 55:209–213
9. Crooks GE, Hon G, Chandonia JM, Brenner SE (2004) WebLogo: a sequence logo generator. *Genome Res* 14:1188–1190
10. Crowther RA, Berriman JA, Curran WL, Allan GM, Todd D (2003) Comparison of the structures of three circoviruses: chicken anemia virus, Porcine circovirus type 2, and Beak and feather disease virus. *Vet Res Commun* 77:13036–13041
11. Dayaram A, Potter KA, Moline AB, Rosenstein DD, Marinov M, Thomas JE, Breitbart M, Rosario K, Arguello-Astorga GR, Varsani A (2013) High global diversity of cycloviruses amongst dragonflies. *J Gen Virol* 94:1827–1840
12. de Villiers E-M, zur Hausen H (2009) TT viruses: the still elusive human pathogens. Springer, Berlin
13. de Villiers EM, Borkosky SS, Kimmel R, Gunst K, Fei JW (2011) The diversity of torque teno viruses: *In vitro* replication leads to the formation of additional replication-competent subviral molecules. *J Virol* 85:7284–7295
14. Delwart E, Li L (2012) Rapidly expanding genetic diversity and host range of the *Circoviridae* viral family and other Rep encoding small circular ssDNA genomes. *Virus Res* 164:114–121
15. Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32:1792–1797
16. Faurez F, Dory D, Grasland B, Jestin A (2009) Replication of porcine circoviruses. *Virol J* 6:60
17. Garigliany M-M, Hagen RM, Frickmann H, May J, Schwarz NG, Perse A, Jöst H, Börstler J, Shahhosseini N, Desmecht D, Mbunkah HA, Daniel AM, Kingsley MT, de Mendonca Campos R, de Paula VS, Randriamampionona N, Poppert S, Tannich E, Rakotozandrindrainy R, Cadar D, Schmidt-Chanasit J (2014) Cyclovirus CyCV-VN species distribution is not limited to Vietnam and extends to Africa. *Sci Rep* 4:7552
18. Ge XY, Li JL, Peng C, Wu LJ, Yang XL, Wu YQ, Zhang YZ, Shi ZL (2011) Genetic diversity of novel circular ssDNA viruses in bats in China. *J Gen Virol* 92:2646–2653
19. Gorbalenya AE, Koonin EV, Wolf YI (1990) A new superfamily of putative NTP-binding domains encoded by genomes of small DNA and RNA viruses. *FEBS Lett* 262:145–148
20. Gorbalenya AE, Koonin EV (1993) Helicases: amino acid sequence comparisons and structure-function relationships. *Curr Opin Struct Biol* 3:419–429
21. Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML 3.0. *Syst Biol* 59:307–321
22. Halami MY, Nieper H, Müller H, John R (2008) Detection of a novel circovirus in mute swans (*Cygnus olor*) by using nested broad-spectrum PCR. *Virus Res* 132:208–212
23. Hamel AL, Lin LL, Nayar GP (1998) Nucleotide sequence of porcine circovirus associated with postweaning multisystemic wasting syndrome in pigs. *J Virol* 72:5262–5267
24. He B, Li Z, Yang F, Zheng J, Feng Y, Guo H, Li Y, Wang Y, Su N, Zhang F, Fan Q, Tu C (2013) Virome profiling of bats from Myanmar by metagenomic analysis of tissue samples reveals more novel mammalian viruses. *PLoS One* 8:e61950
25. Hino S, Prasetyo AA (2009) Relationship of torque teno virus to chicken anemia virus. In: de Villiers E-M, Hausen H (eds) TT viruses: the still elusive human pathogens. Springer, Berlin, pp 117–130
26. Hough KP, Rogers AM, Zelic M, Paris M, Heilman DW (2015) Transformed cell-specific induction of apoptosis by Porcine circovirus type 1 viral protein 3. *J Gen Virol* 96:351–359
27. ICTV (2015) Taxonomy release history. International Committee on Taxonomy of Viruses
28. Ilyina TV, Koonin EV (1992) Conserved sequence motifs in the initiator proteins for rolling circle DNA replication encoded by diverse replicons from eubacteria, eucaryotes and archaeobacteria. *Nucleic Acids Res* 20:3279–3285
29. John R, Fernandez-de-Luaco D, Hofle U, Muller H (2006) Genome of a novel circovirus of starlings, amplified by multiply primed rolling-circle amplification. *J Gen Virol* 87:1189–1195
30. Kamada K, Kuroishi A, Kamahara T, Kabat P, Yamaguchi S, Hino S (2006) Spliced mRNAs detected during the life cycle of Chicken anemia virus. *J Gen Virol* 87:2227–2233
31. Kato A, Fujino M, Nakamura T, Ishihama A, Otaki Y (1995) Gene organization of Chicken anemia virus. *Virology* 209:480–488
32. Kiupel M, Stevenson GW, Galbreath EJ, North A, HogenEsch H, Mittal SK (2005) Porcine circovirus type 2 (PCV2) causes apoptosis in experimentally inoculated BALB/c mice. *BMC Vet Res* 1:1–8
33. Kooistra K, Zhang YH, Henriquez NV, Weiss B, Mumberg D, Noteborn MH (2004) TT virus-derived apoptosis-inducing protein induces apoptosis preferentially in hepatocellular carcinoma-derived cells. *J Gen Virol* 85:1445–1450
34. Li L, Kapoor A, Slikas B, Bamidele OS, Wang C, Shaikat S, Masroor MA, Wilson ML, Ndjanga J-BN, Peeters M, Gross-Camp ND, Muller MN, Hahn BH, Wolfe ND, Triki H, Bartkus J, Zaidi SZ, Delwart E (2010) Multiple diverse circoviruses infect farm animals and are commonly found in human and chimpanzee feces. *J Virol* 84:1674–1682
35. Li L, Victoria JG, Wang C, Jones M, Fellers GM, Kunz TH, Delwart E (2010) Bat guano virome: predominance of dietary viruses from insects and plants plus novel mammalian viruses. *J Virol* 84:6955–6965
36. Li L, Shan TL, Soji OB, Alam MM, Kunz TH, Zaidi SZ, Delwart E (2011) Possible cross-species transmission of circoviruses and cycloviruses among farm animals. *J Gen Virol* 92:768–772
37. Li L, McGraw S, Zhu K, Leutenegger CM, Marks SL, Kubiski S, Gaffney P, Dela Cruz FN Jr., Wang C, Delwart E, Pesavento PA (2013) Circovirus in tissues of dogs with vasculitis and hemorrhage. *Emerg Infect Dis* 19:534–541
38. Li L, Giannitti F, Low J, Keyes C, Ullmann LS, Deng X, Aleman M, Pesavento PA, Pusterla N, Delwart E (2015) Exploring the virome of diseased horses. *J Gen Virol* 96:2721–2733
39. Li L, Pesavento PA, Gaynor AM, Duerr RS, Phan TG, Zhang W, Deng X, Delwart E (2015) A gyrovirus infecting a sea bird. *Arch Virol* 160:2105–2109
40. Li Z, He W, Lan Y, Zhao K, Lv X, Lu H, Ding N, Zhang J, Shi J, Shan C, Gao F (2016) The evidence of porcine hemagglutinating encephalomyelitis virus induced nonsuppurative encephalitis as the cause of death in piglets. *Peer J* 4:e2443
41. Lian H, Liu Y, Li N, Wang Y, Zhang S, Hu R (2014) Novel circovirus from mink, China. *Emerg Infect Dis* 20:1548–1550
42. Lima FE, Cibulski SP, Dall Bello AG, Mayer FQ, Witt AA, Roehe PM, d'Azevedo PA (2015) A novel chiropteran circovirus genome recovered from a Brazilian insectivorous bat species. *Genome Announc* 3:e01393–15
43. Lima FE, Cibulski SP, Dos Santos HF, Teixeira TF, Varela AP, Roehe PM, Delwart E, Franco AC (2015) Genomic

- characterization of novel circular ssDNA viruses from insectivorous bats in Southern Brazil. *PLoS One* 10:e0118070
44. Liu J, Chen I, Kwang J (2005) Characterization of a previously unidentified viral protein in porcine circovirus type 2-infected cells and its role in virus-induced apoptosis. *J Virol* 79:8262–8274
 45. Lőrincz M, Csáagola A, Farkas SL, Széekely C, Tuboly T (2011) First detection and analysis of a fish circovirus. *J Gen Virol* 92:1817–1821
 46. Lőrincz M, Dáan A, Láang M, Csaba G, Tóoth AG, Széekely C, Csáagola A, Tuboly T (2012) Novel circovirus in European catfish (*Silurus glanis*). *Arch Virol* 157:1173–1176
 47. Lukert PD, de Boer GF, Dale JL, Keese P, McNulty MS, Randles JW, Tischer I (1995) Family *Circoviridae*. In: Murphy FA, Fauquet CM, Bishop DHL, Ghabrial SA, Jarvis AW, Martelli GP, Mayo MA, Summers MD (eds) *Virus taxonomy classification and nomenclature of viruses sixth report of the International Committee on Taxonomy of Viruses*. Springer, New York, pp 166–168
 48. Lv Q, Guo K, Zhang G, Zhang Y (2016) The ORF4 protein of porcine circovirus type 2 antagonizes apoptosis by stabilizing the concentration of ferritin heavy chain through physical interaction. *J Gen Virol* 97:1636–1646
 49. Male MF, Kraberger S, Stainton D, Kami V, Varsani A (2016) Cycloviruses, gemycircularviruses and other novel replication-associated protein encoding circular viruses in Pacific flying fox (*Pteropus tonganus*) faeces. *Infect Genet Evol* 39:279–292
 50. Mankertz A, Persson F, Mankertz J, Blaess G, Buhk HJ (1997) Mapping and characterization of the origin of DNA replication of porcine circovirus. *J Virol* 71:2562–2566
 51. Mankertz A, Mankertz J, Wolf K, Buhk HJ (1998) Identification of a protein essential for replication of porcine circovirus. *J Gen Virol* 79:381–384
 52. Mankertz A, Hattermann K, Ehlers B, Soike D (2000) Cloning and sequencing of columbid circovirus (CoCV), a new circovirus from pigeons. *Arch Virol* 145:2469–2479
 53. Meehan BM, Creelan JL, McNulty MS, Todd D (1997) Sequence of porcine circovirus DNA: affinities with plant circoviruses. *J Gen Virol* 78:221–227
 54. Miyata H, Tsunoda H, Kazi A, Yamada A, Khan MA, Murakami J, Kamahora T, Shiraki K, Hino S (1999) Identification of a novel GC-rich 113-nucleotide region to complete the circular, single-stranded DNA genome of TT virus, the first human circovirus. *J Virol* 73:3582–3586
 55. Muhire B, Martin DP, Brown JK, Navas-Castillo J, Moriones E, Zerbini FM, Rivera-Bustamante R, Malathi VG, Briddon RW, Varsani A (2013) A genome-wide pairwise-identity-based proposal for the classification of viruses in the genus *Mastrevirus* (family *Geminiviridae*). *Arch Virol* 158:1411–1424
 56. Muhire BM, Varsani A, Martin DP (2014) SDT: a virus classification tool based on pairwise sequence alignment and identity calculation. *PLoS One* 9:e108277
 57. Niagro FD, Forsthoefel AN, Lawther RP, Kamalanathan L, Ritchie BW, Latimer KS, Lukert PD (1998) Beak and feather disease virus and porcine circovirus genomes: intermediates between the geminiviruses and plant circoviruses. *Arch Virol* 143:1723–1744
 58. Noteborn MHM, Koch G (1995) Chicken anemia virus infection: molecular basis of pathogenicity. *Avian Pathol* 24:11–31
 59. Padilla-Rodriguez M, Rosario K, Breitbart M (2013) Novel cyclovirus discovered in the Florida woods cockroach *Eurycotis floridana* (Walker). *Arch Virol* 158:1389–1392
 60. Peters MA, Jackson DC, Crabb BS, Browning GF (2002) Chicken anemia virus VP2 is a novel dual specificity protein phosphatase. *J Biol Chem* 277:39566–39573
 61. Phan TG, Li L, O’Ryan MG, Cortes H, Mamani N, Bonkougou IJO, Wang C, Leutenegger CM, Delwart E (2012) A third gyrovirus species in human faeces. *J Gen Virol* 93:1356–1361
 62. Phan TG, Phung Vo N, Sdiri-Loulizi K, Aouni M, Pothier P, Ambert-Balay K, Deng X, Delwart E (2013) Divergent gyroviruses in the feces of Tunisian children. *Virology* 446:346–348
 63. Phan TG, Luchsinger V, Avendano LF, Deng X, Delwart E (2014) Cyclovirus in nasopharyngeal aspirates of Chilean children with respiratory infections. *J Gen Virol* 95:922–927
 64. Phan TG, Mori D, Deng X, Rajindrajith S, Ranawaka U, Fan Ng TF, Bucardo-Rivera F, Orlandi P, Ahmed K, Delwart E (2015) Small circular single stranded DNA viral genomes in unexplained cases of human encephalitis, diarrhea, and in untreated sewage. *Virology* 482:98–104
 65. Prasetyo AA, Kamahora T, Kuroishi A, Murakami K, Hino S (2009) Replication of chicken anemia virus (CAV) requires apoptin and is complemented by VP3 of human torque teno virus (TTV). *Virology* 385:85–92
 66. Rijsewijk FAM, dos Santos HF, Teixeira TF, Cibulski SP, Varela APM, Dezen D, Franco AC, Roehle PM (2011) Discovery of a genome of a distant relative of chicken anemia virus reveals a new member of the genus Gyrovirus. *Arch Virol* 156:1097–1100
 67. Rinder M, Schmitz A, Peschel A, Korbel R (2015) Complete genome sequence of a novel circovirus from zebra finch. *Genome Announc* 3:e00560–15
 68. Ritchie BW, Niagro FD, Lukert PD, Steffens WL, Latimer KS (1989) Characterization of a new virus from cockatoos with psittacine beak and feather disease. *Virology* 171:83–88
 69. Ritchie BW, Niagro FD, Latimer KS, Lukert PD, Steffens WL 3rd, Rakich PM, Pritchard N (1990) Ultrastructural, protein composition, and antigenic comparison of psittacine beak and feather disease virus purified from four genera of psittacine birds. *J Wildl Dis* 26:196–203
 70. Rosario K, Marinov M, Stainton D, Kraberger S, Wiltshire EJ, Collings DA, Walters M, Martin DP, Breitbart M, Varsani A (2011) Dragonfly cyclovirus, a novel single-stranded DNA virus discovered in dragonflies (Odonata: Anisoptera). *J Gen Virol* 92:1302–1308
 71. Rosario K, Dayaram A, Marinov M, Ware J, Kraberger S, Stainton D, Breitbart M, Varsani A (2012) Diverse circular ssDNA viruses discovered in dragonflies (Odonata: Epiprocta). *J Gen Virol* 93:2668–2681
 72. Rosario K, Duffy S, Breitbart M (2012) A field guide to eukaryotic circular single-stranded DNA viruses: insights gained from metagenomics. *Arch Virol* 157:1851–1871
 73. Sato G, Kawashima T, Kiuchi M, Tohya Y (2015) Novel cyclovirus detected in the intestinal contents of Taiwan squirrels (*Callosciurus erythraeus taiwanensis*). *Virus Genes* 51:148–151
 74. Sauvage V, Cheval J, Foulongne V, Gouilh MA, Pariente K, Manuguerra JC, Richardson J, Dereure O, Lecuit M, Burguiere A, Caro V, Eloit M (2011) Identification of the first human *Gyrovirus*, a virus related to chicken anemia virus. *J Virol* 85:7948–7950
 75. Simmonds PM, AMJ, Benkő M, Breitbart M, Brister JR, Carstens EB, Davison AJ, Delwart E, Gorbalenya AE, Harrach B, Hull R, King AMQ, Koonin EV, Krupovic M, Kuhn JH, Lefkowitz EJ, Nibert ML, Orton R, Roossinck MJ, Sabanadzovic S, Sullivan MB, Suttle CA, Tesh RB, van der Vlugt RA, Varsani A, Zerbini FM (in press) *Virus taxonomy in the age of metagenomics*. *Nature Rev Microbiol*. doi:10.1038/nrmicro.2016.1177
 76. Smits SL, Zijlstra EE, van Hellemond JJ, Schapendonk CM, Bodewes R, Schurch AC, Haagmans BL, Osterhaus AD (2013) Novel cyclovirus in human cerebrospinal fluid, Malawi, 2010–2011. *Emerg Infect Dis* 19:1511–1513
 77. Steinfeldt T, Finsterbusch T, Mankertz A (2006) Demonstration of nicking/joining activity at the origin of DNA replication

- associated with the Rep and Rep' proteins of porcine circovirus type 1. *J Virol* 80:6225–6234
78. Stewart ME, Perry R, Raidal SR (2006) Identification of a novel circovirus in Australian ravens (*Corvus coronoides*) with feather disease. *Avian Pathol* 35:86–92
 79. Takahashi K, Iwasa Y, Hijikata M, Mishiro S (2000) Identification of a new human DNA virus (TTV-like mini virus, TLMV) intermediately related to TT virus and chicken anemia virus. *Arch Virol* 145:979–993
 80. Tan le V, van Doorn HR, Nghia HD, Chau TT, Tu le TP, de Vries M, Canuti M, Deijis M, Jebbink MF, Baker S, Bryant JE, Tham NT, NT BK, Boni MF, Loi TQ, Phuong le T, Verhoeven JT, Crusat M, Jeeninga RE, Schultsz C, Chau NV, Hien TT, van der Hoek L, Farrar J, de Jong MD (2013) Identification of a new cyclovirus in cerebrospinal fluid of patients with acute central nervous system infections. *mBio* 4:e00231–00213
 81. Todd D, Creelan JL, Mackie DP, Rixon F, McNulty MS (1990) Purification and biochemical characterization of chicken anemia agent. *J Gen Virol* 71:819–823
 82. Todd D, Niagro FD, Ritchie BW, Curran W, Allan GM, Lukert PD, Latimer KS, Steffens WL 3rd, McNulty MS (1991) Comparison of three animal viruses with circular single-stranded DNA genomes. *Arch Virol* 117:129–135
 83. Todd D (2000) Circoviruses: immunosuppressive threats to avian species: a review. *Avian Pathol* 29:373–394
 84. Todd D, Weston J, Ball NW, Borghmans BJ, Smyth JA, Gelmini L, Lavazza A (2001) Nucleotide sequence-based identification of a novel circovirus of canaries. *Avian Pathol* 30:321–325
 85. Todd D, Weston JH, Soike D, Smyth JA (2001) Genome sequence determinations and analyses of novel circoviruses from goose and pigeon. *Virology* 286:354–362
 86. Todd D, Scott ANJ, Fringuelli E, Shivraprasad HL, Gavier-Widen D, Smyth JA (2007) Molecular characterization of novel circoviruses from finch and gull. *Avian Pathol* 36:75–81
 87. Varsani A, Martin DP, Navas-Castillo J, Moriones E, Hernandez-Zepeda C, Idris A, Zerbini FM, Brown JK (2014) Revisiting the classification of curtoviruses based on genome-wide pairwise identity. *Arch Virol* 159:1873–1882
 88. Varsani A, Navas-Castillo J, Moriones E, Hernandez-Zepeda C, Idris A, Brown JK, Murilo Zerbini F, Martin DP (2014) Establishment of three new genera in the family *Geminiviridae*: *Becurtovirus*, *Eragrovirus* and *Turncurtovirus*. *Arch Virol* 159:2193–2203
 89. Varsani A, Krupovic M (in press) Sequence-based taxonomic framework for the classification of uncultured single-stranded DNA viruses of the family *Genomoviridae*. *Virus Evol*
 90. Wu Z, Ren X, Yang L, Hu Y, Yang J, He G, Zhang J, Dong J, Sun L, Du J, Liu L, Xue Y, Wang J, Yang F, Zhang S, Jin Q (2012) Virome analysis for identification of novel mammalian viruses in bat species from Chinese provinces. *J Virol* 86:10999–11012
 91. Wu Z, Yang L, Ren X, He G, Zhang J, Yang J, Qian Z, Dong J, Sun L, Zhu Y, Du J, Yang F, Zhang S, Jin Q (2016) Deciphering the bat virome catalog to better understand the ecological diversity of bat viruses and the bat origin of emerging infectious diseases. *ISME J* 10:609–620
 92. Zhang W, Li L, Deng X, Kapusinszky B, Pesavento PA, Delwart E (2014) Faecal virome of cats in an animal shelter. *J Gen Virol* 95:2553–2564