N-Methyl-D-aspartate and α -amino-3-hydroxy-5-methyl-4-isoxazolepropionate receptors involved in the induction of sedative effects under an acute stress in neonatal chicks

**H. Yamane · Y. Tsuneyoshi · D. M. Denbow ·
M. Furuse**

Published online: 31 March 2009
© Springer-Verlag 2009

Erratum to: Amino Acids (2009)
DOI 10.1007/s00726-008-0203-x

In the published original version of this article, first page, Abstract, 15th line, the information regarding the potency of NMDA (“less”) is not given correct.

The correct sentence should read:

Although NMDA also induced a sedative effect, the potency of NMDA for sleep-like behavior was less than that of glutamate.

The online version of the original article can be found under doi:[10.1007/s00726-008-0203-x](https://doi.org/10.1007/s00726-008-0203-x).

H. Yamane · Y. Tsuneyoshi · M. Furuse (✉)
Laboratory of Advanced Animal and Marine Bioresources,
Graduate School of Bioresources and Bioenvironmental
Sciences, Kyushu University, Fukuoka 812-8581, Japan
e-mail: furuse@brs.kyushu-u.ac.jp

D. M. Denbow
Department of Animal and Poultry Sciences,
Virginia Polytechnic Institute and State University,
Blacksburg, VA 24061-0306, USA