Correction to: Cone beam computed tomography and ultrasonography imaging of benign intraosseous jaw lesion: a prospective radiopathological study

Ibrahim Sevki Bayrakdar¹ · Ahmet Berhan Yilmaz² · Fatma Caglayan² · Umit Ertas³ · Cemal Gundogdu⁴ · Ismail Gumussoy⁵

Published online: 8 February 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Correction to: Clinical Oral Investigations
<https://doi.org/10.1007/s00784-017-2257-1>

In the original version of this article, ‘123 lytic intraosseous jaw lesions diagnosed in 112 patients’ was incorrectly presented as ‘123 lytic intraosseous jaw lesions diagnosed in 121 patients’ and ‘an average age of 31.7 ± 15.4 (range, 6–72)’ was incorrectly presented as ‘average age of 15.4 ± 31.7 (range, 6–72)’.

The online version of the original article can be found at <https://doi.org/10.1007/s00784-017-2257-1>

✉ Ibrahim Sevki Bayrakdar
ibrahimsevkibayrakdar@gmail.com

- ¹ Department of Dentomaxillofacial Radiology, Faculty of Dentistry, Eskisehir Osmangazi University, Eskisehir, Turkey
- ² Department of Dentomaxillofacial Radiology, Faculty of Dentistry, Ataturk University, Erzurum, Turkey
- ³ Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Ataturk University, Erzurum, Turkey
- ⁴ Medical Pathology, Erzurum, Turkey
- ⁵ Turkish Health Ministry Gungoren Oral Health Hospital, Istanbul, Turkey