Microbial diversity and methanogenic potential in a high temperature natural gas field in Japan

Hanako Mochimaru · Hideyoshi Yoshioka · Hideyuki Tamaki ·
Kohei Nakamura · Nobuyuki Kaneko · Susumu Sakata · Hiroyuki Imachi ·
Yuji Sekiguchi · Hiroo Uchiyama · Yoichi Kamagata

Published online: 24 May 2008
© Springer 2008

Erratum to: Extremophiles (2007) 11:453–461
DOI 10.1007/s00792-006-0056-8

The unit of acetate concentration in Table 1 and the reference in the text was shown incorrectly. The unit should be mmol l^{-1} . This does not affect any of our conclusions.

The online version of the original article can be found under doi:[10.1007/s00792-006-0056-8](https://doi.org/10.1007/s00792-006-0056-8).

H. Mochimaru · H. Uchiyama
Graduate School of Life and Environmental Sciences,
University of Tsukuba, Tsukuba,
Ibaraki 305-8572, Japan

H. Mochimaru · H. Tamaki · K. Nakamura ·
Y. Sekiguchi · Y. Kamagata (✉)
Institute for Biological Resources and Functions,
National Institute of Advanced Industrial Science
and Technology (AIST), 1-1-1 Higashi, Tsukuba,
Ibaraki 305-8566, Japan
e-mail: y.kamagata@aist.go.jp

Y. Kamagata
Research Institute of Genome-Based Biofactory,
National Institute of Advanced Industrial Science
and Technology (AIST), Sapporo, Hokkaido 062-8517, Japan

H. Yoshioka · N. Kaneko · S. Sakata
Institute for Geo-resources and Environment,
National Institute of Advanced Industrial Science
and Technology (AIST), Tsukuba, Ibaraki 305-8566, Japan

H. Imachi
Subground Animalcule Retrieval (SUGAR) Program,
Extremobiosphere Research Center,
Japan Agency for Marine-Earth Science & Technology
(JAMSTEC), Yokosuka, Kanagawa 237-0061, Japan