Silvia Bagnoli · Andrea Tedde · Elena Cellini ·
Mario Rotondi · Benedetta Nacmias · Sandro Sorbi

The urokinase-plasminogen activator (PLAU) gene is not associated with late onset Alzheimer's disease

Published online: 19 March 2005
© Springer-Verlag 2005

Neurogenetics (2004) 6:53–54

Table 1 was omitted owing to technical problems. It is now printed below.

Table 1 PLAUI P141L polymorphism: genotypes and corresponding allele frequencies in AD and controls

Study	Sample/ size	Genotype			Allele		Association	
		TT %	TC %	CC %	T% %	C% %	Genotypic	Allelic
Bagnoli 2004	AD (238)	2.1	26	71.8	15.1	84.8	$P>0.1$	$P>0.1$
	CTR (207)	2.9	24.1	72.9	14.9	85	$\chi^2=0.462$; 2 <i>df</i>	$\chi^2=0.00$; 1 <i>df</i>
Myers 2004	AD (652)	7	46	73	24	76	$P>0.1$	$P>0.1$
	CTR (824)	6	37	58	24	76	$\chi^2=0.1$; 2 <i>df</i>	$\chi^2=0.02$; 1 <i>df</i>
Finckh 2003	AD (347)	4	28.5	67.5	17.2	81.7	$P>0.00039$	$P>0.00009$
	CTR (291)	7.2	40.6	52.2	27.4	72.5	$\chi^2=15.689$; 2 <i>df</i>	$\chi^2=15.341$; 1 <i>df</i>

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10048-004-0203-2>

S. Bagnoli (✉) · A. Tedde · E. Cellini ·
B. Nacmias · S. Sorbi
Department of Neurological and Psychiatric Sciences,
University of Florence,
Viale Pieraccini 6,
50139 Florence, Italy
e-mail: bagnolis@unifi.it
Tel.: +39-055-4271379
Fax: +39-055-4271380

M. Rotondi
Department of Clinical Pathophysiology,
Unit of Endocrinology University of Florence,
Florence, Italy