The levels of sCD30 and of sCD40L in a group of patients with systemic lupus erythematoses and their diagnostic value

Hana Ciferská · Pavel Horák · Zuzana Heřmanová ·
Marta Ordeltová · Josef Zadražil · Tomáš Tichý ·
Vlastimil Ščudla

Published online: 2 March 2007
© Clinical Rheumatology 2007

Clin Rheumatol
DOI 10.1007/s10067-006-0389-9

Figure 2 describes the levels of sCD40L. The label on the y-axis incorrectly refers to SCD30.

The legend and the figure themselves are correct.

The Czech text in the first paragraph of “Discussion” appeared inadvertently and should have been deleted.

Fig. 2 The levels of sCD40L in the group with SLE in the healthy controls and in the subgroups with lupus nephritis and without lupus nephritis and with the high (ECLAM>3) and low activity of SLE (ECLAM≤3). Statistical difference was found between the whole group of patients with SLE and the healthy controls ($p=0.02$)

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10067-006-0389-9>.

H. Ciferská · P. Horák · J. Zadražil · V. Ščudla
III. Department of Internal Medicine, Faculty of Medicine
and Faculty Hospital, University of Olomouc,
Olomouc, Czech Republic

Z. Heřmanová · M. Ordeltová
Department of Clinical Immunology, Faculty Hospital,
Olomouc, Czech Republic

T. Tichý
Department of Pathology, Faculty of Medicine
and Faculty Hospital, University of Olomouc,
Olomouc, Czech Republic

H. Ciferská (✉)
III. interní klinika FNO,
I.P. Pavlova,
6775 20 Olomouc, Czech Republic
e-mail: hana.ciferska@volny.cz