ERRATUM

Erratum to: Predictors of poor sleep quality in patients with systemic lupus erythematosus

M. Inoue¹ · K. Shiozawa² · R. Yoshihara² · T. Yamane³ · Y. Shima⁴ · T. Hirano⁴ · K. Makimoto¹

Published online: 3 May 2017

© International League of Associations for Rheumatology (ILAR) 2017

Erratum to: Clin Rheumatol
DOI 10.1007/s10067-017-3545-5

The original version of this article, unfortunately, contained errors.

In Fig. 1, three out of four *p* values and the names of categories were missing. The corrected image is shown in this article.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10067-017-3545-5>

✉ K. Makimoto
kmakimot@sahs.med.osaka-u.ac.jp

¹ Department of Nursing, Graduate School of Medicine, Osaka University, 1-7 Yamadaoka, Suita, Osaka 565-0871, Japan

² Rheumatology & Collagen Disease Center, Hyogo Prefectural Kakogawa Medical Center, 203 Kanno, Kanno-cho, Kakogawa, Hyogo 675-0003, Japan

³ Rheumatology Center, Kakogawa City Hospital, 439 Honmachi, Kakogawa-cho, Kakogawa, Hyogo 675-0037, Japan

⁴ Department of Allergy and Rheumatic Disease, Graduate School of Medicine, Osaka University, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan

Fig. 1 Comparison between good sleeper and poor sleeper on SF-12 and LupusPRO summary score *ES* effect size, *HRQoL* LupusPRO health related quality of life, *Non-Health QoL* LupusPRO non-health related quality of life, *MCS* short form health survey-12 mental component summary score, *PCS* short form health survey-12 physical component summary score