ERRATUM

K. Masumoto · I. Yamada · H. Tanaka · Y. Fujise
K. Hashimoto

Tissue distribution of a new photosensitizer ATX-S10Na(II) and effect of a diode laser (670 nm) in photodynamic therapy

Published online: 23 January 2004
© Springer-Verlag London Limited 2004

Lasers Med Sci (2003) 18:134–138

Mistakenly, all authors were listed with the same affiliation. Their correct affiliations are given here.

In the last sentence of the section Material and methods p. 135, “ig/g” should have been “μg/g”.

Table 1 was reproduced incorrectly. The correct table is given here.

Table 1 The correlation between irradiation condition and antitumor effect (+ depth of necrosis ≥ 1.0 mm, < 1.0 mm)

mW/cm ² /s	30	60	100	200	250	313	400	500	625	800	1000	1500
Group 1: ATX-S10Na(II) + irradiation												
300	– (n=3)	+/- (n=2/1)	+ (n=3)	+ (n=1)								
200		– (n=1)	– (n=3)	+ (n=3)	+ (n=3)	+ (n=1)						
160			– (n=3)	+/- (n=1/2)	+ (n=3)		+ (n=3)		+ (n=3)		+ (n=3)	
100				– (n=3)	– (n=3)	+/- (n=2/1)	+ (n=3)	+ (n=3)				
60						– (n=3)	+/- (n=1/2)	+ (n=3)	+ (n=1)			
Group 2: Irradiation only												
300	– (n=1)	– (n=2)	– (n=3)	+/- (n=2/1)	+ (n=3)		+ (n=2)					
200					– (n=1)	– (n=3)	+/- (n=2/1)	+ (n=3)	+ (n=1)			
160									– (n=2)	– (n=2)	– (n=3)	
100												
60												

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10103-003-0268-4>

K. Masumoto (✉) · H. Tanaka
Department of Oral and Maxillofacial Surgery,
Hamamatsu University School of Medicine,
1-20-1 Handayama, Hamamatsu, 431-3192 Japan
E-mail: kazusan@me.ndmc.ac.jp

I. Yamada · K. Hashimoto
Department of Endoscopic and Photodynamic Medicine,
Hamamatsu University School of Medicine,
1-20-1 Handayama, Hamamatsu, 431-3192 Japan

Y. Fujise
Department of Chemistry,
Hamamatsu University School of Medicine,
1-20-1 Handayama, Hamamatsu, 431-3192 Japan