

Heat conduction from the exceedingly hot fiber tip contributes to the endovenous laser ablation of varicose veins

Renate R. van den Bos · Michael A. Kockaert ·
H. A. Martino Neumann · Rolf H. Bremmer ·
Tamar Nijsten · Martin J. C. van Gemert

Published online: 4 April 2009
© Springer-Verlag London Limited 2009

Erratum to: Lasers Med Sci
DOI 10.1007/s10103-008-0639-y

Unfortunately Fig. 3 was omitted and Fig. 2 was repeated in its place.

The correct Fig. 3 is reproduced here.




Fig. 3 As Fig. 2, now for a vein radius of $r=2.5$ mm and $T_{tip}=1,000^{\circ}\text{C}$

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10103-008-0639-y>.

R. R. van den Bos · M. A. Kockaert · H. A. Martino Neumann ·
T. Nijsten (✉)
Department of Dermatology, Erasmus Medical Centre,
P.O. BOX 2040, 3000 CA Rotterdam, The Netherlands
e-mail: t.nijsten@erasmusmc.nl

R. H. Bremmer · M. J. C. van Gemert
Department of Biomedical Engineering and Physics,
Academic Medical Centre,
Amsterdam, The Netherlands