Erratum to: EURO-CORDEX: new high-resolution climate change projections for European impact research

Daniela Jacob · Juliane Petersen · Bastian Eggert · Antoinette Alias · Ole Bøssing Christensen · Laurens M. Bouwer · Alain Braun · Augustin Colette · Michel Déqué · Goran Georgievski · Elena Georgopoulou · Andreas Gobiet · Laurent Menut · Grigory Nikulin · Andreas Haensler · Nils Hempelmann · Colin Jones · Klaus Keuler · Sari Kovats · Nico Kröner · Sven Kotlarski · Arne Kriegsmann · Eric Martin · Erik van Meijgaard · Christopher Moseley · Susanne Pfeifer · Swantje Preuschmann · Christine Radermacher · Kai Radtke · Diana Rechid · Mark Rounsevell · Patrick Samuelsson · Samuel Somot · Jean-Francois Soussana · Claas Teichmann · Riccardo Valentini · Robert Vautard · Björn Weber · Pascal Yiou

Published online: 12 March 2014
© Springer-Verlag Berlin Heidelberg 2014

Erratum to: Reg Environ Change DOI 10.1007/s10113-013-0499-2

The author would like to correct the errors in the publication of the original article. The corrected details are given below for your reading.

In result section, third sentence of the fourth paragraph should read as:

“The zone between regions in which precipitation increases in the north and decreases in the south shifts northwards in summer and southwards in winter”

The online version of the original article can be found under doi:[10.1007/s10113-013-0499-2](https://doi.org/10.1007/s10113-013-0499-2).

D. Jacob (✉) · J. Petersen · B. Eggert · A. Haensler · N. Hempelmann · A. Kriegsmann · C. Moseley · S. Pfeifer · S. Preuschmann · D. Rechid · C. Teichmann · B. Weber
Climate Service Center (CSC), Helmholtz-Zentrum Geesthacht, Fischertwiete 1, 20095 Hamburg, Germany
e-mail: daniela.jacob@hzg.de

D. Jacob · J. Petersen · C. Moseley · C. Radermacher · D. Rechid · C. Teichmann
Max Planck Institute for Meteorology (MPIM), Bundesstr. 53, 20146 Hamburg, Germany

A. Alias · A. Braun · M. Déqué · E. Martin · S. Somot
Météo-France/CNRS, Centre National de Recherches Météorologiques (CNRM-GAME), 42, Avenue Gaspard Coriolis, 31057 Toulouse Cedex 1, France

O. B. Christensen
Danish Climate Centre, Danish Meteorological Institute (DMI), Lyngbyvej 100, 2100 Copenhagen, Denmark

L. M. Bouwer
Institute for Environmental Studies, Vrije Universiteit, De Boelelaan 1087, 1081 HV Amsterdam, The Netherlands

L. M. Bouwer
Deltares, PO Box 177, 2600 MH Delft, The Netherlands

A. Colette
Institut National de l'Environnement Industriel et des Risques (INERIS), 60550 Verneuil-en-Halatte, France

G. Georgievski · K. Keuler · K. Radtke
Environmental Meteorology, Brandenburg University of Technology, Burger Chaussee 2, 03013 Cottbus, Germany

E. Georgopoulou
Institute for Environmental Research and Sustainable Development (IERSD), National Observatory of Athens, 118 10 Thessio, Greece

A. Gobiet
Wegener Center for Climate and Global Change and Institute for Geophysics, Astrophysics, and Meteorology, University of Graz, Brandhofgasse 5, 8010 Graz, Austria

L. Menut
Laboratoire de Météorologie Dynamique (IPSL), Ecole Polytechnique, 91128 Palaiseau Cedex, France

The ranges in Table 3 should be:

Cold spell duration index, Max, Continental, RCP8.5 from -6 to -8 .

Annual total precipitation, Min, Southern, RCP4.5 from -11 to 2 .

Annual total precipitation, Max, Southern, RCP4.5 from 2 to -11 .

The correct Table 3 is given in the following page.

G. Nikulin · C. Jones · P. Samuelsson
Swedish Meteorological and Hydrological Institute (SMHI),
Folkborgsvägen 17, 601 76 Norrköping, Sweden

S. Kovats
London School of Hygiene and Tropical Medicine,
15-17 Tavistock Place, London WC1H 9 SH, UK

N. Kröner · S. Kotlarski
Institute for Atmospheric and Climate Science, ETH Zurich,
Universitätsstrasse 16, 8092 Zurich, Switzerland

E. van Meijgaard
Royal Netherlands Meteorological Institute (KNMI),
Wilhelminalaan 10, 3732 GK De Bilt, The Netherlands

M. Rounsevell
David Kinloch Michie Chair of Rural Economy &
Environmental Sustainability, Institute of Geography & the
Lived Environment, School of GeoSciences, The University of
Edinburgh, Drummond Street, Edinburgh EH8 9XP, UK

J.-F. Soussana
INRA, UR874, Grassland Ecosystems Research (UREP),
Clermont-Ferrand, France

R. Valentini
Department of Forest Science and Environment,
University of Tuscia, Viterbo, Italy

R. Valentini
Euro-Mediterranean Center for Climate Change, Via De Lellis,
01100 Viterbo, Italy

R. Vautard · P. Yiou
Laboratoire des Sciences du Climat et de l'Environnement,
IPSL, CEA/CNRS/UVSQ, Gif sur Yvette, France

Table 3 Projected changes of selected climate parameters and indices for 2071–2100 with respect to 1971–2000 spatially averaged for European sub-regions based on RCP4.5 and RCP8.5

Scenarios	Climate parameters	Measure	Alpine			Atlantic			Continental			Northern			Southern		
			RCP 4.5	RCP 8.5	RCP 4.5	RCP 8.5	RCP 4.5	RCP 8.5	RCP 4.5	RCP 8.5	RCP 4.5	RCP 8.5	RCP 4.5	RCP 8.5	RCP 4.5	RCP 8.5	RCP 8.5
2071–2100 minus 1971–2000	Mean annual temperature in K	Median	2.4	4.6	1.7	3.2	2.1	4.1	2.9	5.2	2.0	4.2	2.0	5.2	2.0	4.2	4.2
		Min	1.8	3.8	1.3	2.5	1.6	3.6	2.0	4.1	1.9	3.8	1.9	4.1	1.9	3.8	3.8
		Likely in the range	1.9–3.4	3.9–6.0	1.4–2.1	2.7–3.6	1.6–3.2	3.7–5.2	2.0–4.2	4.1–6.2	1.9–2.7	3.9–5.4	1.9–2.7	4.1–6.2	1.9–2.7	3.9–5.4	3.9–5.4
	Frost days (1) per year	Max	3.6	6.3	2.9	4.2	3.2	5.3	4.3	6.5	3.2	5.7	3.2	6.5	3.2	5.7	5.7
		Median	–40	–70	–28	–40	–34	–62	–40	–68	–22	–43	–22	–68	–22	–43	–43
		Min	–25	–55	–12	–21	–16	–46	–24	–58	–10	–22	–10	–58	–10	–22	–22
	Likely in the range	Likely in the range	–26 to –41	–57 to –85	–15 to –30	–26 to –50	–18 to –40	–50 to –65	–26 to –43	–60 to –83	–11 to –30	–23 to –51	–11 to –30	–60 to –83	–11 to –30	–23 to –51	–23 to –51
		Max	–47	–93	–33	–60	–41	–73	–52	–93	–31	–51	–31	–93	–31	–51	–51
		Median	8	19	11	24	20	37	4	13	27	54	4	13	27	54	54
	Summer days (2) per year	Min	3	10	6	17	11	27	2	5	21	43	2	5	21	43	43
		Likely in the range	4–14	12–24	6–14	22–28	13–24	30–46	2–16	6–22	25–33	46–60	2–16	6–22	25–33	46–60	46–60
	Max	Max	18	25	33	38	28	49	23	28	37	67	23	28	37	67	67
		Median	1	4	3	7	9	22	1	1	20	45	1	1	20	45	45
		Min	0	1	0	3	2	11	0	0	7	23	0	0	7	23	23
	Likely in the range	Likely in the range	1–3	2–5	1–5	3–12	9–27	17–31	0–5	1–3	11–24	25–57	0–5	1–3	11–24	25–57	25–57
		Max	8	6	18	17	30	37	7	13	41	58	7	13	41	58	58
		Median	31	61	39	58	26	58	23	55	27	49	23	55	27	49	49
	Min	Min	23	52	24	41	17	52	17	37	16	34	17	37	16	34	34
		Likely in the range	23–39	52–83	27–43	47–68	20–38	53–71	19–33	41–60	17–33	38–53	19–33	41–60	17–33	38–53	38–53
		Max	45	95	45	75	41	75	42	78	38	58	42	78	38	58	58
	Warm spell duration index (14) days per year	Median	34	96	20	65	23	73	35	82	34	124	35	82	34	124	124
		Min	26	73	17	46	16	52	22	64	28	90	22	64	28	90	90
		Likely in the range	29–55	77–136	20–31	49–87	18–42	58–93	23–42	75–113	32–69	98–177	23–42	75–113	32–69	98–177	98–177
	Max	Max	69	162	55	102	54	106	63	130	83	186	63	130	83	186	186
		Median	–5	–5	–5	–5	–6	–6	–7	–6	–5	–5	–7	–6	–5	–5	–5
		Min	–3	–3	–2	–4	–3	–5	–4	–4	–3	–4	–4	–4	–3	–4	–4
	Likely in the range	Likely in the range	–4 to –7	–4 to –6	–3 to –6	–4 to –5	–4 to –7	–5 to –8	–6 to –8	–5 to –7	–3 to –5	–4 to –5	–6 to –8	–5 to –7	–3 to –5	–4 to –5	–4 to –5
		Max	–7	–6	–6	–6	–7	–8	–8	–8	–6	–6	–8	–8	–6	–6	–6
		Median	5	14	1	4	9	10	10	22	–6	–10	10	22	–6	–10	–10
	Annual total precipitation (27) in %	Min	3	5	–1	–2	0	0	7	18	2	0	7	18	2	0	0
		Likely in the range	4 to 8	7 to 16	–1 to 7	2 to 9	1 to 13	4 to 19	8 to 17	19 to 32	–10 to 0	–21 to –2	8 to 17	19 to 32	–10 to 0	–21 to –2	–21 to –2
		Max	12	18	9	9	16	29	22	34	–11	–27	22	34	–11	–27	–27
	Annual total precipitation where RR >99p of 1971/2000 (26) in %	Median	38	79	36	71	44	65	43	82	36	49	43	82	36	49	49
		Min	24	41	20	48	17	37	27	64	23	30	27	64	23	30	30
		Likely in the range	25–60	44–105	25–70	49–105	33–60	44–75	28–65	66–110	31–57	38–58	28–65	66–110	31–57	38–58	38–58
	Max	Max	73	119	73	118	73	106	70	120	62	65	70	120	62	65	65

Numbers are based on 8 (RCP4.5) and 9 (RCP8.5) regional model simulations. The likely range defines the range of 66 % of all projected changes around the ensemble median