Proteomics of Juvenile Senegal Sole (*Solea senegalensis*) Affected by Gas Bubble Disease in Hyperoxygenated Ponds

E. Salas-Leiton · B. Cánovas-Conesa · R. Zerolo ·
J. López-Barea · J. P. Cañavate · J. Alhama

Published online: 6 May 2009
© Springer Science + Business Media, LLC 2009

Erratum to: Mar Biotechnol
DOI 10.1007/s10126-008-9168-8

In the original version of this article, there were some errors on the citation of figure components in section “Identification of Differentially Expressed Proteins” and in Figs. 3 and 4 legends.

The second sentence of section “Identification of Differentially Expressed Proteins” should read:

In gill, only RKIP (spot 8) was most intense in GBD-affected animals without symptoms, while the other seven identified proteins, β -ACT (spot 3), RKIP (spot 5), CaM, β -TBB, α -TPM/MLC1, β -ACT (spot 22), and GAPDH, were most intense in affected fish with symptoms, most of them following expression pattern I.

Correct Figs. 3 and 4 legends are given here.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10126-008-9168-8>.

E. Salas-Leiton · R. Zerolo · J. P. Cañavate
IFAPA Centro El Toruño, Junta de Andalucía,
Apartado 16,
11500 El Puerto de Santa María, Cádiz, Spain

B. Cánovas-Conesa · J. López-Barea · J. Alhama (✉)
Departamento de Bioquímica y Biología Molecular,
Universidad de Córdoba,
Edificio S. Ochoa,
14071 Córdoba, Spain
e-mail: bb2alcaj@uco.es

Fig. 3 1 Master gel of gill cytosolic proteins with 25 spots characteristic of six expression types: (A) present only and (B) absent in healthy fish. (E) Increased and (F) diminished in affected asymptomatic fish. (G) Present only in affected fish with external symptoms. (I) Increased in affected fish with external symptoms. 2 Relative intensities of the proteins included in the master gel in the three statuses studied: healthy control fish; GBD-affected asymptomatic animals; GBD-affected fish with visible symptoms

Fig. 4 1 Master gel of liver cytosolic proteins with 23 spots characteristic of eight expression types: (A) present, (B) absent, and (C) increased in healthy fish. (D) Absent and (F) diminished in affected asymptomatic fish. (G) Present, (H) absent, and (I) increased in affected fish with external symptoms. 2 Relative intensities of proteins included in the master gel in the three health statuses studied: healthy control fish; GBD-affected asymptomatic animals; GBD-affected fish with visible symptoms