Erratum to: Risk-stratified survival rates and predictors of biochemical recurrence after radical prostatectomy in a Nara, Japan, cohort study

Nobumichi Tanaka · Kiyohide Fujimoto · Akihide Hirayama · Kazumasa Torimoto ·
Eijiro Okajima · Masahiro Tanaka · Makito Miyake · Keiji Shimada ·
Noboru Konishi · Yoshihiko Hirao

Published online: 9 February 2013
© Japan Society of Clinical Oncology 2013

Erratum to: Int J Clin Oncol (2011) 16:553–559
DOI 10.1007/s10147-011-0226-2

Part of Table 1 (rows 29–42 of the original) was incorrectly shown. The correct data are given here.

Table 1 (partial)

	Low risk (<i>n</i> = 122)	Intermediate risk (<i>n</i> = 131)	High risk (<i>n</i> = 215)	<i>p</i> value
ECE				
Yes	26	40	78	0.017*
No	96	91	137	
PNI				
Yes	44	52	95	ns*
No	72	65	89	
Unknown	6	14	31	
SVI				
Yes	5	3	31	<0.001*
No	117	128	184	
PSM				
Yes	24	25	55	ns*
No	85	87	125	
Unknown	13	19	35	

The online version of the original article can be found under doi:[10.1007/s10147-011-0226-2](https://doi.org/10.1007/s10147-011-0226-2).

N. Tanaka (✉) · K. Fujimoto · A. Hirayama · K. Torimoto ·
Y. Hirao
Department of Urology, Nara Medical University,
840 Shijo-cho, Kashihara, Nara 634-8522, Japan
e-mail: sendo@naramed-u.ac.jp

E. Okajima · M. Tanaka · M. Miyake
Nara Uro-Oncological Research Group,
840 Shijo-cho, Kashihara, Nara 634-8522, Japan

K. Shimada · N. Konishi
Department of Pathology, Nara Medical University,
840 Shijo-cho, Kashihara, Nara 634-8522, Japan

ECE external capsular extension, *PNI* perineural invasion, *SVI* seminal vesicle involvement, *PSM* positive surgical margin, *ns* not significant

* Chi-squared test