Infralevator leiomyoma

A. Kamrava · M. A. Abbas · D. S. Katz

Received: 2 March 2009 / Accepted: 9 October 2010 / Published online: 18 November 2010
© Springer-Verlag 2010

A 39-year-old woman presented with chronic and cyclical anorectal pain. A right ischiorectal mass was palpable but there were no skin changes. Ultrasound examination showed a solid mass extending from within the levator plate (Fig. 1). Computed tomography confirmed a $5.7 \times 3.2 \times 5.0$ cm mass (Fig. 2). The patient underwent a pararectal excision of the mass (Fig. 3). On histological

Fig. 1 Ultrasound examination showing a solid mass extending from within the levator plate

Fig. 2 Computed tomography confirmed a $5.7 \times 3.2 \times 5.0$ cm mass

Fig. 3 Intraoperative image of the tumor protruding from the anal verge

A. Kamrava (✉) · M. A. Abbas
Kaiser Permanente Hospital, Los Angeles, CA, USA
e-mail: allen.kamrava@gmail.com

D. S. Katz
Winthrop-University Hospital, New York, NY, USA

Fig. 4 Specimen of the pararectal excision of the mass revealing a leiomyoma

examination revealed a leiomyoma (Fig. 4). The inferior aspect of the levator ani muscle is an uncommon location for a leiomyoma.