Effects of high sodium intake and diuretics on the circadian rhythm of blood pressure in type 2 diabetic patients treated with an angiotensin II receptor blocker

Takashi Uzu · Masayoshi Sakaguchi · Yukiyo Yokomaku · Shinji Kume · Masami Kanasaki · Keiji Isshiki · Shin-ichi Araki · Toshiro Sugimoto · Daisuke Koya · Masakazu Haneda · Atsunori Kashiwagi

Published online: 14 April 2009
© Japanese Society of Nephrology 2009

Erratum to: Clin Exp Nephrol DOI 10.1007/s10157-009-0157-7

The legend for Fig. 3 appeared incorrectly in the article cited above. The correct legend is as follows.

The online version of the original article can be found under doi:[10.1007/s10157-009-0157-7](https://doi.org/10.1007/s10157-009-0157-7).

T. Uzu (✉) · M. Sakaguchi · Y. Yokomaku · S. Kume · M. Kanasaki · K. Isshiki · S. Araki · T. Sugimoto · A. Kashiwagi
Department of Medicine, Shiga University of Medical Science,
Otsu, Shiga, Japan
e-mail: takuzu@belle.shiga-med.ac.jp

D. Koya
Division of Endocrinology and Metabolism,
Department of Medicine, Kanazawa Medical University,
Kanazawa, Japan

M. Haneda
Department of Medicine, Asahikawa Medical College,
Asahikawa, Japan

Fig. 3 Mean change in BP values from baseline in 24-h mean, daytime, night-time and morning SBP and DBP obtained after 24 weeks of treatment with losartan (50 mg) plus hydrochlorothiazide (12.5 mg) (white bars) and valsartan monotherapy (160 mg) (black bars). Mean $\pm$ SD, $\dagger P < 0.05$ and $*P < 0.01$ between treatments. SBP systolic blood pressure, DBP diastolic blood pressure