Erratum to: Prevalence of chronic kidney disease in the Japanese general population

Enyu Imai · Masaru Horio · Tsuyoshi Watanabe · Kunitoshi Iseki · Kunihiro Yamagata · Shigeko Hara · Nobuyuki Ura · Yutaka Kiyohara · Toshiki Moriyama · Yasuhiro Ando · Shoichi Fujimoto · Tsuneo Konta · Hitoshi Yokoyama · Hirofumi Makino · Akira Hishida · Seiichi Matsuo

Published online: 22 October 2009
© Japanese Society of Nephrology 2009

Erratum to: Clin Exp Nephrol DOI 10.1007/s10157-009-0199-x

In Table 3, in the column headed “Proteinuria (+)”, the “Estimated number of Japanese adults in 2005” in the

30–59 age-group should be 823881, not 823881. The corrected table is shown here.

The online version of the original article can be found under doi: [10.1007/s10157-009-0199-x](https://doi.org/10.1007/s10157-009-0199-x).

E. Imai (✉)
Department of Nephrology,
Osaka University Graduate School of Medicine,
Suita, Osaka 565-0871, Japan
e-mail: ADS12069@nifty.com

M. Horio
Department of Functional Diagnostic Science,
Osaka University Graduate School of Medicine,
Osaka, Japan

T. Watanabe
Third Department of Medicine,
Fukushima Medical University, Fukushima, Japan

K. Iseki
Dialysis Unit, University Hospital of the Ryukyus,
Nishihara, Okinawa, Japan

K. Yamagata
Department of Nephrology, Institute of Clinical Medicine,
Graduate School of Comprehensive Human Sciences,
University of Tsukuba, Ibaraki, Japan

S. Hara
Health Medical Center, Toranomon Hospital,
Tokyo, Japan

N. Ura
Department of General Medicine,
Teine Keijinkai Hospital, Sapporo, Hokkaido, Japan

Y. Kiyohara
Department of Environmental Medicine, Graduate School
of Medical Sciences, Kyushu University, Fukuoka, Japan

T. Moriyama
Healthcare Center, Osaka University, Osaka, Japan

Y. Ando
Department of Nephrology, Jichi Medical School,
Tochigi, Japan

S. Fujimoto
First Department of Medicine, Miyazaki University,
Miyazaki, Japan

T. Konta
First Department of Medicine, Yamagata University,
Yamagata, Japan

H. Yokoyama
Division of Nephrology, Kanazwa Medical University,
Ishikawa, Japan

H. Makino
Department of Nephrology, Diabetes and Rheumatology,
Okayama University Graduate School of Medicine,
Dentistry and Pharmaceutical Sciences, Okayama, Japan

A. Hishida
First Department of Medicine, Hamamatsu University School of
Medicine, Shizuoka, Japan

Table 3 Prevalence rates of CKD stages in Japanese adults (20 years or older), and estimated number of CKD cases per CKD stage based on the 2005 census

GFR (ml/min/1.73 m ²)	Total	Proteinuria (+)	Proteinuria (–)
Prevalence rate (%)			
GFR $\geq$ 90	27.8	0.6	27.2
60–89	61.6	1.7	60.0
30–59	10.4	0.8	9.6
<30	0.2	0.1	0.1
Stage 3			
50–59	7.6	0.4	7.2
40–49	2.3	0.3	2.0
30–39	0.6	0.1	0.4
Estimated number of Japanese adults in 2005			
GFR $\geq$ 90	28639274	605313	28033961
60–89	63576938	1708870	61868068
30–59	10743236	823881	9919355
<30	236569	125190	111379
Stage 3			
50–59	7809261	425146	7384116
40–49	2363987	267158	2096828
30–39	569988	131577	438411