Erratum to: Chronic Kidney Disease Japan Cohort study: baseline characteristics and factors associated with causative diseases and renal function

Enyu Imai · Seiichi Matsuo · Hirofumi Makino ·
Tsuyoshi Watanabe · Tadao Akizawa · Kosaku Nitta ·
Satoshi Iimuro · Yasuo Ohashi · Akira Hishida

Published online: 6 October 2010
© Japanese Society of Nephrology 2010

Erratum to: Clin Exp Nephrol
DOI 10.1007/s10157-010-0328-6

In the Appendix, the name of Daijo Inaguma was misspelled in the original version as Daijyo Inaguma.

In Tables 1 and 4, the numbers given for Creatinine clearance values were incorrect. The correct values are indicated on the following page.

For The CKD-JAC Study Group.

The online version of the original article can be found under doi:[10.1007/s10157-010-0328-6](https://doi.org/10.1007/s10157-010-0328-6).

E. Imai (✉) · S. Matsuo
Department of Nephrology, Nagoya University Graduate School of Medicine, 65 Tsurumai, Showa-ku, Nagoya 466-8550, Japan
e-mail: imai@med.nagoya-u.ac.jp

H. Makino
Department of Medicine and Clinical Science,
Okayama University Graduate School of Medicine,
Dentistry and Pharmaceutical Sciences, Okayama, Japan

T. Watanabe
The Third Department of Internal Medicine,
Fukushima Medical College, Fukushima, Japan

T. Akizawa
Department of Nephrology, Showa University, Tokyo, Japan

K. Nitta
Department of Medicine, Kidney Center,
Tokyo Women's Medical University, Tokyo, Japan

S. Iimuro · Y. Ohashi
Department of Biostatistics, School of Public Health,
The University of Tokyo, Tokyo, Japan

A. Hishida
First Department of Medicine, Hamamatsu University School of Medicine, Hamamatsu, Japan

Table 1 Patient characteristics classified by causative disease

Variable	Cohort, <i>N</i> = 2977	No diabetes		Diabetes		<i>P</i> value
		No CGN, <i>N</i> = 909	CGN, <i>N</i> = 948	No nephropathy, <i>N</i> = 507	Nephropathy, <i>N</i> = 613	
Ccr (ml/min)						
Mean (SD)	48.03 (29.98)	44.31 (26.34)	49.12 (30.05)	51.36 (32.83)	48.74 (32.20)	0.1095
Median (max–min)	41.70 (4.8–240.0)	39.62 (7.0–151.7)	44.10 (4.8–240.0)	42.95 (10.7–172.5)	41.80 (11.7–180.3)	
1Q–3Q	26.90–59.50	24.80–57.50	27.80–59.70	29.30–59.69	24.50–62.10	

Table 4 Baseline characterization

	Stage 3A GFR $\geq$ 45, <i>N</i> = 304	Stage 3B 45 > GFR $\geq$ 30, <i>N</i> = 1037	Stage 4 30 > GFR $\geq$ 15, <i>N</i> = 1160	Stage 5 GFR < 15, <i>N</i> = 476	<i>P</i> value
Ccr (ml/min)					
Mean (SD)	90.65 (34.77)	62.88 (27.75)	37.78 (15.37)	20.92 (9.11)	<0.0001
Median (min–max)	82.05 (30.9–180.3)	56.40 (8.8–240.0)	34.31 (7.2–97.7)	19.30 (4.8–53.8)	
1Q–3Q	67.25–114.09	45.20–71.20	26.85–46.10	14.81–25.09	