Erratum to: U-shaped association between body mass index and proteinuria in a large Japanese general population sample

Yuji Sato · Shouichi Fujimoto · Tsuneo Konta · Kunitoshi Iseki · Toshiki Moriyama ·
Kunihiro Yamagata · Kazuhiko Tsuruya · Hideaki Yoshida · Koichi Asahi ·
Issei Kurahashi · Yasuo Ohashi · Tsuyoshi Watanabe

Published online: 28 June 2013
© Japanese Society of Nephrology 2013

Erratum to: Clin Exp Nephrol DOI 10.1007/s10157-013-0809-5

The original version of this article unfortunately contained errors.

In the Abstract, under the heading “Methods”, the number of men (median age 66 years) should be 85,183, not 185,183.

Also in the “Methods” section, under the heading “Baseline measurement”, lines 11–14 should read:

Urine dipstick results were interpreted by the medical staff at each local medical institution and recorded as (–), (±), (1±), (2±), and (3±).

The online version of the original article can be found under doi:[10.1007/s10157-013-0809-5](https://doi.org/10.1007/s10157-013-0809-5).

Y. Sato (✉)
Dialysis Division, University of Miyazaki Hospital,
Miyazaki, Japan
e-mail: ysato@fc.miyazaki-u.ac.jp

S. Fujimoto
Department of Hemovascular Medicine and Artificial Organs,
Faculty of Medicine, University of Miyazaki, Miyazaki, Japan

T. Konta
Department of Cardiology, Pulmonology, and Nephrology,
Yamagata University School of Medicine, Yamagata, Japan

K. Iseki · T. Moriyama · K. Yamagata · K. Tsuruya ·
H. Yoshida · K. Asahi · T. Watanabe
Steering Committee for the “Examination of the Positioning
of CKD in Specific Health Check and Guidance”, Tokyo, Japan

I. Kurahashi
Department of Planning, Information, and Management,
University of Tokyo Hospital, Tokyo, Japan

Y. Ohashi
Department of Biostatistics/Epidemiology and Preventive Health
Sciences, School of Health Sciences and Nursing, University
of Tokyo, Tokyo, Japan