ERRATUM

Erratum to: Impact of hemoglobin levels on renal and non-renal clinical outcomes differs by chronic kidney disease stages: the Gonryo study

Tae Yamamoto¹ · Mariko Miyazaki^{1,2} · Masaaki Nakayama^{2,3} · Gen Yamada¹ · Masato Matsushima^{2,4} · Mistuhiro Sato⁵ · Toshinobu Sato⁵ · Yoshio Taguma^{2,5} · Hiroshi Sato^{2,6} · Sadayoshi Ito^{1,2}

Published online: 26 June 2017
© Japanese Society of Nephrology 2017

Erratum to: Clin Exp Nephrol (2016) 20:595–602 DOI 10.1007/s10157-015-1190-3

In the original publication Fig. 1 was incorrectly published. The revised version of Fig. 1 is given in this Erratum.

The online version of the original article can be found under doi:[10.1007/s10157-015-1190-3](https://doi.org/10.1007/s10157-015-1190-3).

✉ Tae Yamamoto
tae.yamamoto@med.tohoku.ac.jp

¹ Division of Nephrology, Endocrinology and Vascular Medicine, Tohoku University Graduate School of Medicine, 1-1 Seiryomachi, Aoba-ku, Sendai, Miyagi 980-8574, Japan

² Center for Advanced Integrated Renal Science, Tohoku University Graduate School of Medicine, Sendai, Japan

³ Division of Nephrology, Hypertension, Diabetology, Endocrinology, and Metabolism, Fukushima Medical University School of Medicine, Fukushima, Japan

⁴ Department of Clinical Research, The Jikei University School of Medicine, Tokyo, Japan

⁵ Kidney Center, Japan Community Health Care Organization Sendai Hospital, Sendai, Japan

⁶ Clinical Pharmacology and Therapeutics, Tohoku University Graduate School of Pharmaceutical Sciences, Sendai, Japan