Altered ivermectin pharmacology and defective visual system in *Drosophila* mutants for histamine receptor HCLB

Shazie Yusein · Nadya Velikova · Petia Kупenova ·
Roger Hardie · Adrian Wolstenholme ·
Eugene Semenov

Published online: 21 November 2008
© Springer-Verlag 2008

Erratum to: Invert Neurosci
DOI 10.1007/s10158-008-0078-2

The original version of this article unfortunately contained a mistake. The presentation of Fig. 6 was published with an error. The correct version is given on the next page.

The online version of the original article can be found under
doi:[10.1007/s10158-008-0078-2](https://doi.org/10.1007/s10158-008-0078-2).

S. Yusein · N. Velikova · P. Kупenova · E. Semenov (✉)
Department of Molecular Neurobiology,
Institute of Molecular Biology, 1113 Sofia, Bulgaria
e-mail: esen@obzor.bio21.bas.bg

R. Hardie
Department of Physiology, Development and Neuroscience,
Cambridge University, Cambridge CB2 3DY, UK

A. Wolstenholme
Department of Biology and Biochemistry, University of Bath,
Claverton Down, Bath BA2 7AY, UK

Fig. 6 Allele-specific changes of the transient components of the ERGs following ivermectin application. The number of responses registered under specific experimental conditions for a given experimental group of 10 animals is shown in Table 3. **a, d** and **g** *OR/Df(3R)E79*; **b, e** and **h** *hclB^{T1}/Df(3R)E79*; **c, f** and **i** *hclB^{T2}/Df(3R)E79*