Erratum

Erratum to: Antioxidants Reduce Cellular and Functional Changes Induced by Intense Noise in the Inner Ear and Cochlear Nucleus

JIANZHONG LU,¹ WEI LI,¹ XIAOPING DU,¹ DONALD L. EWERT,¹ MATTHEW B. WEST,¹ CHARLES STEWART,² ROBERT A. FLOYD,² AND RICHARD D KOPKE^{1,2,3}

¹*Hough Ear Institute, 3400 N.W. 56th Street, Oklahoma City, OK 73112, USA*

²*Oklahoma Medical Research Foundation, Oklahoma City, OK 73104, USA*

³*Departments of Physiology and Otolaryngology, University of Oklahoma Health Sciences Center, Oklahoma City, OK 73104, USA*

Erratum to: JARO

DOI 10.1007/s10162-014-0441-4

The labels for the Y axis in Figure 10 were incorrectly labeled “% Missing OHCs” and should read “% Missing IHCs” as shown here.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10162-014-0441-4>.

Correspondence to: Richard D Kopke · Hough Ear Institute · 3400 N.W. 56th Street, Oklahoma City, OK 73112, USA.
rkopke@houghear.org

FIG. 10. Percent IHC loss as a function of percent distance (frequency place in kilohertz) from the OC apex in normal control (no noise, untreated) and at 4 h (A), 12 h (B), 24 h (C), 7 days (D), and 21 days (E) after exposure to the 115-dB OBN in the noise only and treated groups. Shaded areas show the frequency range of the noise. The data are plotted as mean±SEM. Six cochlea were analyzed for

each data point. IHC loss was first seen in the basal part of the cochlea 4 h post-noise exposure and increased and expanded over the 21 days after noise exposure. Significant differences were seen at 4 h, 12 h, 7 days, and 21 days after noise exposure. Data analysis was performed by two-way ANOVA followed by Bonferroni post-tests: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.