Erratum to: The ratio of transmitted near-infrared radiation to photosynthetically active radiation (PAR) increases in proportion to the adsorbed PAR in the canopy

Atsushi Kume · Kenlo N. Nasahara ·
Shin Nagai · Hiroyuki Muraoka

Published online: 9 July 2010
© The Botanical Society of Japan and Springer 2010

Erratum to: J Plant Res
DOI 10.1007/s10265-010-0346-1

In Fig. 4 some symbols (circles) in part **a** were incorrectly shown. In the legend for part **b**, a minus sign (−) was used in the equation; it should have been a plus sign (+). The corrected figure for **a** and legend for **b** appear below:

The online version of the original article can be found under doi:10.1007/s10265-010-0346-1.

A. Kume (✉)
Ashoro Research Forest, Department of Forest and Forest
Products Sciences, Faculty of Agriculture, Kyushu University,
1-85 Kita-5jo, Ashoro, Hokkaido 089-3705, Japan
e-mail: akume@attglobal.net

K. N. Nasahara
Graduate School of Life and Environmental Science,
University of Tsukuba, 1-1-1 Tennodai,
Tsukuba 305-8572, Japan

S. Nagai
Research Institute for Global Change,
Japan Agency for Marine-Earth Science and Technology,
3173-25 Showa-machi, Kanazawa-ku,
Yokohama 236-0001, Japan

H. Muraoka
Institute for Basin Ecosystem Studies, Gifu University,
1-1 Yanagido, Gifu 501-1193, Japan

Fig. 4 Relationship between NIRt/PArt and **a** PARt/PArt ($y = 14.48x - 11.52$; $R^2 = 0.96$; $n = 13$), and **b** LAI [$y = 2.80 \ln(x) + 0.69$; $R^2 = 0.97$; $n = 13$]