Erratum to: Fate and transport of bensulfuron-methyl and imazosulfuron in paddy fields: experiments and model simulation

Kazuhiro Takagi · Ferdinand F. Fajardo ·
Masumi Ishizaka · Thai Khanh Phong ·
Hirozumi Watanabe · Julien Boulange

Published online: 22 July 2011
© Springer-Verlag 2011

Erratum to: Paddy Water Environ
DOI 10.1007/s10333-011-0276-0

Due to an unfortunate turn of events this article has been published with an erroneous version of Figs. 4, 5 and 6. Please find in this erratum the correct Figs. 4, 5 and 6 that should be regarded by the reader as the final version.

The online version of the original article can be found under doi:[10.1007/s10333-011-0276-0](https://doi.org/10.1007/s10333-011-0276-0).

K. Takagi (✉) · M. Ishizaka
National Institute for Agro-Environmental Sciences, Tsukuba,
Ibaraki 305-8604, Japan
e-mail: ktakagi@niaes.affrc.go.jp

F. F. Fajardo
De La Salle University, 2401 Taft Avenue, Malate 1004, Manila,
Philippines

T. K. Phong
Queensland University, 39 Kessels Road, Coopers Plains,
QLD 4108, Australia

H. Watanabe · J. Boulange
Tokyo University of Agriculture and Technology, 3-5-8
Saiwaicho, Fuchu, Tokyo 183-8509, Japan

Fig. 4 Simulated (*continuous line*) and observed (*open square*) bensulfuron-methyl (BSM) concentrations in paddy water (*above*) and soil paddy field (*below*)

Fig. 5 Simulated (*continuous line*) and observed (*open square*) imazosulfuron (IMS) concentrations in paddy water (*above*) and soil paddy field (*below*)

Fig. 6 Pesticide mass balance as % applied mass for bensulfuron-methyl (*above*) and imazosulfuron (*below*)