



Correction to: Simultaneous Quantification of 5-Aminoimidazole-4-Carboxamide-1- β -D-ribofuranoside and Its Active Metabolite 5-Aminoimidazole-4-Carboxamide-1- β -D-ribofuranotide in Mice Plasma by LC–MS/MS

Xiaoliang Cheng¹ · Bing Chen² · Yuanyuan Pan² · Liping Guo² · Weiyi Feng¹ · Yalin Dong¹

Published online: 24 November 2017
© Springer-Verlag GmbH Germany, part of Springer Nature 2017

Correction to: Chromatographia
<https://doi.org/10.1007/s10337-017-3392-1>

In the original version of this article the presentation of Figs 1, 2, 3 was incorrect.

The corrected figures are given below.




Fig. 1 Chemical structure of 5-aminoimidazole-4-carboxamide-1- β -D-ribofuranoside (a) and 5-aminoimidazole-4-carboxamide-1- β -D-ribofuranotide (b)

The original article can be found online at <https://doi.org/10.1007/s10337-017-3392-1>.

✉ Weiyi Feng
fengweiyi@hotmail.com

✉ Yalin Dong
dongyalin@mail.xjtu.edu.cn

¹ Department of Pharmacy, The First Affiliated Hospital of Xi'an Jiaotong University, Xi'an 710061, China

² Thermo Fisher Scientific (China) Co., Ltd, Shanghai, China




Fig. 2 Full-scan mass spectra for the parent and product of AICA riboside and AICA ribotide: **a** parent ion spectrum for AICA riboside, **b** parent ion spectrum for AICA ribotide, **c** product ion spectrum for AICA riboside, and **d** product ion spectrum for AICA ribotide




Fig. 3 Multiple reaction monitoring chromatograms of AICA riboside and its active metabolite AICA ribotide in mice plasma: **a** blank mice plasma, **b** blank mice plasma spiked with 1 µg/mL AICA riboside, 1 µg/mL AICA ribotide, or internal standard; **c** AICA ribo-

side and AICA ribotide in a plasma sample obtained from a mice at 15 min following a single intravenous dose (200 mg/kg) of AICA riboside