Multimodal encoding in a simplified model of intracellular calcium signaling

Maurizio De Pittà · Vladislav Volman ·
Herbert Levine · Eshel Ben-Jacob

Published online: 10 January 2009
© Marta Olivetti Belardinelli and Springer-Verlag 2008

Erratum to: Cogn Process
DOI 10.1007/s10339-008-0242-y

The legend of Fig. 1 contains errors. Please find below
Fig. 1 with the correct figure legend.

Fig. 1 **a** Purified astrocytes from rat's visual cortex loaded with calcium indicator fluo-3 and stimulated by norepinephrine (25 μ M) and **(b)** associated normalized Ca^{2+} traces. Similarly to astrocytes, many other cells display intracellular Ca^{2+} oscillations in response to several external agents. Such oscillations are characterized by great variability both in their peaks and in their frequency, hinting complex modes of encoding of information of external signals. Images and experimental data courtesy of V. Parpura [adapted from: Lee W, Parpura V (2007) Exocytotic release of glutamate from astrocytes: comparison to neurons. In: Bean A (ed) Protein trafficking in neurons. Elsevier, Amsterdam, pp 329–365]

The online version of the original article can be found under
doi:[10.1007/s10339-008-0242-y](https://doi.org/10.1007/s10339-008-0242-y).

M. De Pittà · E. Ben-Jacob (✉)
School of Physics and Astronomy, Tel Aviv University,
69978 Ramat Aviv, Israel
e-mail: eshel@tamar.tau.ac.il

V. Volman · H. Levine · E. Ben-Jacob
Center for Theoretical Biological Physics,
UCSD, La Jolla, CA 92093-0319, USA

V. Volman
Computational Neurobiology Lab, The Salk Institute,
La Jolla, CA 92037, USA